



Antimicrobial Activity of Chitosan Film Forming Solution Enriched with Essential Oils; an *in Vitro* Assay

Kana Jean Raphaël¹ and Amir Meimandipour^{*2}

¹Department of Animal Productions, Faculty of Agronomy and Agricultural Sciences, University of Dschang, 70 Dschang, Cameroon

²Department of Animal Biotechnology, National Institute of Genetic Engineering and Biotechnology, Tehran, 14965/161, Iran

*Corresponding author: Amir Meimandipour, Department of Animal Biotechnology, National Institute of Genetic Engineering and Biotechnology, Tehran, Iran. Tel: +98-2144787459, Fax: +98-2144787395, E-mail: meimandi@nigeb.ac.ir

Received: 25 Oct. 2015; Revised: 2 July 2016; Accepted: 19 June 2017; Published online: 19 August 2017

Background: The resistance of the bacteria and fungi to the innumerable antimicrobial agents is a major challenge in the treatment of the infections demands to the necessity for searching and finding new sources of substances with antimicrobial properties. The incorporation of the essential oils (EOs) in chitosan film forming solution may enhance antimicrobial properties. However, its use as the feeding additive in the poultry nutrition needs to clarify the product's activity against both pathogen and the useful microbes in the gastrointestinal tract.

Objectives: In the present study, we carried out an *in vitro* investigation and evaluated the antimicrobial activity of chitosan film forming solution incorporated with essential oils (CFs+EOs) against microbial strains including *Staphylococcus aureus*, *Escherichia coli*, *Enterococcus faecium*, *Lactobacillus rahnmosus*, *Aspergillus niger* and *Alternaria alternate*.

Material and Methods: In three replicates, the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of different treatments including: 1- essential oils (EOs), 2- chitosan film solution (CFs), and 3- chitosan film solution enriched with EOs (CFs+EOs) were determined against above mentioned microbes.

Results: The results indicated that the chitosan solution enriched with essential oils (CFs+EOs) is capable of inhibiting the bacterial and fungal growth even at the lowest concentrations. The MIC and MBC for all the antimicrobial agents against *Escherichia coli* and *Staphylococcus aureus* were very low compared to the concentrations needed to inhibit the growth of useful bacteria, *Lactobacillus rahnmosus* and *Enterococcus faecium*. The antifungal activity of chitosan was enhanced as the concentration of EOs increased in the film solution.

Conclusion: Chitosan-EOs complexes are the promising candidate for novel contact antimicrobial agents that can be used in animal feeds.

Keywords: Antimicrobial properties; Bacteria; Chitosan, Chitosan film forming solution; Essential oils; Fungi

1. Background

There are many probiotics that emerged into the market that have the potential to increase the growth performances of the chickens. These probiotics include spices, plant extracts, and essential oils (EOs). Essential oils are volatile compounds that are found in aromatic plants all over the world. They are sources of biologically active compounds with strong antibacterial, antioxidant, and digestion stimulating abilities (1-2). The use of these volatile compounds as feed additives (3-7) has become an attractive area of research in poultry science. Essential oils from a variety of the plants have been shown to stimulate feed intake

and digestion, improve body weight gain, support of a positive gut microflora, and offer healthy performance benefits to poultry (8-9). The enhancement of the animal growth performance by EOs is due to the reduction of pathogens within the intestine with direct consequence of increasing nutrient availability for animal utilization due to the lower nutrient competition and intestinal diseases prevention (1). The other beneficial effects of the supplementing animal feed with EOs include stimulation of appetite, improvement of digestive enzyme secretion, immune system modulation, antimicrobial activity and decreased mortality (7, 10-13). As well, there is a synergism when several EOs

are mixed, the result of which is a greater antimicrobial effect than individual EOs alone (1-2, 14).

Due to their oxidative and volatile properties in addition to their susceptibility to high temperatures, EOs have a tendency to lose their efficiency due to the mitigated quality. The main components of EOs are labile, volatile, and most of them are easily lost during feed processing and storage. Under different storage conditions and environmental variation, EOs degrade rapidly when compared to chemotherapies. The volatile property of EOs causes a large loss in the quantity of the product during feed processing and storage. Incorporation in a stable matrix can overcome the technical issues of stability and odour in the feed (15). Moreover, incorporation in a matrix can stabilize and avoid loss of EOs quality and quantity in animal feed formulation. It may also lead to flavor optimization of the feed, better handling, reduce dustiness and increase stability, delay in the release of EOs in the digestive tract, and enhancement of bioavailability (16). In this regard, hydrophilic nanoparticles of chitosan having the capacity to bind varieties of biomolecules have recently attracted attentions in the human food industry.

Chitosan is a natural biodegradable and biocompatible non-toxic polymer extracted from crustacean shells having the potential to efficiently retain the bioactivity of molecules such as nucleic acids, proteins, and antigens (17-21). In addition, it also presents immune-stimulating, antimicrobial, wound-healing properties (22-25). In addition, it has recently been used as the vaccine delivery systems (26-27) as well as volatile feed additives compounds such as EOs in animal feed (15). Their antimicrobial property is effective only in aqueous systems and this property becomes inefficient in its insoluble film aspect (28). In the later aspect, chitosan constitutes a barrier capable of reducing respiration and retards bacterial and fungal growth (29-30). According to Ruiz-Navajas *et al.* (33), chitosan film solutions enriched with EOs enhance the chitosan antimicrobial properties and stabilize lipid in the feed. Up to now, few studies have been reported about chitosan-compounded fragrance for animal feed. Mixing chitosan with EOs could allow them to exert their *in vivo* antimicrobial and digestive-stimulating activities by ensuring a more efficient delivery to the target site in the gastrointestinal tract.

2. Objectives

The present *in vitro* study was proposed to give an overview on the potential of chitosan film for retaining and protecting the essential oil bioactivity and for

evaluating the benefits of the association chitosan-essential oil toward supporting a positive gut microfloral growth.

3. Materials and Methods

3.1. Materials

Medium molecular weight chitosan [Poly (D-glucosamine)] was provided from Sigma-Aldrich (St. Louis, MO, USA). MRS broth, Nutrient broth, and Sabouraud Dextrose broth were procured from Iranian Bioresearch CO. All reagents used in this study were of analytical grade. Oregano and thyme essential oils were procured from Barij Esans Co. Iran. In the experiment, all solutions were prepared with distilled water.

3.2. Preparation of Chitosan Film (CFs) and Chitosan Film Incorporated with Essential Oils- (CFs+EOs) Solutions

Chitosan films solution (CFs) were prepared according to Chi *et al.* (28) with modification. Chitosan stock solution (1% (w/v)) was prepared under magnetic stirring by dissolving 1 g of chitosan in 100 mL of an aqueous acetic acid solution (1% (v/v)) at ambient temperature overnight. A blend of Oregano (500 μ L) and Thyme (500 μ L) EOs was mixed with 500 μ L of tween 20 and introduced in the chitosan stock solution under constant stirring. The final essential oils-loaded chitosan film forming solution consisted of 1% chitosan, 1% acetic acid, 0.5% Tween 20, and 1% EOs. The freshly prepared CFs and CF+EOs solutions were used for antimicrobial assays.

3.3. Structure of Chitosan Films

FTIR spectra of chitosan powder, chitosan films (CFs), EOs-loaded chitosan film (CFs+EOs), Thyme EO, Oregano EO and blend of Thyme and Oregano EOs were recorded to evaluate the cross-linking of chitosan with EOs in the film. FTIR spectra were recorded on a BRUKER FTIR spectrophotometer (SENSOR 27, Germany) using potassium bromide (KBr) pellets at a resolution of 4 cm^{-1} .

3.4. Preparation of Antimicrobial Essential Oil Solution

A blend of the EOs solution was prepared by mixing 2.5 mL of Oregano and 2.5 mL of Thyme essential oils in 2.5 mL of tween 20. Then, the mixture of EOs-tween 20 was added to 100 mL distilled water under magnetic stirring for 30 min and used for antimicrobial assays. CFs and EOs solutions were used as control samples for testing the antimicrobial activity of CFs+EOs. The concentrations of EOs in the solution was tested

ranging from 50 to 0.39 $\mu\text{L}\cdot\text{mL}^{-1}$ (50, 25, 12.5, 6.25, 3.12, 1.56, 0.78 and 0.39 $\mu\text{L}\cdot\text{mL}^{-1}$) while in CFs+EOs solution, EOs concentrations was ranged from 10 to 0.078 $\mu\text{L}\cdot\text{mL}^{-1}$ (10, 5, 2.5, 1.25, 0.625, 0.312, 0.156 and 0.078 $\mu\text{L}\cdot\text{mL}^{-1}$).

3.5. Microbial Strains

Inhibition assays were carried out with pure cultures of the bacterial and fungal strains. A total of four different bacteria were used; three grams of the positive and one gram of the negative genera. The Gram positive strains were *Enterococcus faecium* (PTCC 1237), *Lactobacillus rahnmosus* (PTCC 1637) and *Staphylococcus aureus* (PTCC 1189). A Gram negative bacterium was *Escherichia coli* (PTCC 1399). To prepare the stock cultures, *Enterococcus faecium* and *Lactobacillus rahnmosus* were grown in MRS agar and *Staphylococcus aureus* and *Escherichia coli* were grown on nutrient agar and were stored under refrigeration condition at 4-8 °C and subcultured weekly. Two molds, *Aspergillus niger* (PTCC 5154) and *Alternaria alternata* (PTCC 5224) were also tested. Cultures were grown in Sabouraud Dextrose Agar and stored under refrigerated conditions at 4-8 °C and subcultured weekly. All test micro-organisms were obtained from the culture collection of the National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran.

3.6. Antimicrobial Activity Assessment

The minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of CFs, CFs+EOs and EOs solutions were determined as described by Qi *et al.* (22). Briefly, a number of test tubes, each containing 5.0 mL of MRS broth as culture media for *Enterococcus faecium* and *Lactobacillus rahnmosus*, and nutrient broth for *Escherichia coli* and *Staphylococcus aureus* were autoclaved for 15 min at 121 °C. To the first tube of each, 5.0 mL of CFs, CFs+EOs or EOs solutions as prepared above was respectively added. After mixing, 5.0 mL of the mixture was transferred to the second tube, and similar transformations were repeated. Each tube contained a test sample solution with half of the concentration of the previous one. The tubes were inoculated under aseptic conditions with 50 μL (1×10^7 CFU. mL^{-1}) of the freshly prepared bacterial suspension and incubated in an orbital shaker (180 rpm) at 37 °C for 20 h. The control tubes were contained only MRS broth and nutrient broth. After incubation, a loopful from each tube was cultured on its relevant culture medium and incubated at 37 °C for 24 h.

The MIC was defined as the lowest concentration of compound that completely inhibited visible growth of bacteria. The MBC was defined as the lowest concentration of CFs, CFs+EOs, and EOs that kills all the bacteria, resulting in no growth on subculture.

3.7. Antifungal Activity Assay

Chitosan film (CFs), chitosan film enriched with EOs (CFs+EOs) and pure essential oils (EOs) solutions were screened for antifungal activity against *Aspergillus niger* and *Alternaria alternata* by observing the percentage inhibition of mycelial growth under their effect according to a method modified from the procedure reported by Kaur *et al.* (25). About 20 mL of the Sabouraud Dextrose Agar (SDA) medium was poured into Petri plates and allowed to solidify. After solidification, 1 mL of CFs, CFs+EOs, and EOs stock solution were spread on top using a sterilized swab and were dried for 10 min. 5 mm discs of a 7-days-old culture of the test fungi were placed at the center of the above Petri plates and incubated at 37 °C for 96 h. After incubation, the diameter (mm) of the grown colony was measured. For each treatment, three replicates were maintained. Sabouraud Dextrose Agar medium without the CFs, CFs+EOs, and EOs solution served as control. The minimal inhibitory concentration (MIC) of each compound was defined as the lowest concentration of compound that completely inhibited visible growth after 96 h of incubation. The minimal fungicidal concentration (MFC) was defined as the lowest concentration resulting in no growth on subculture.

The toxicity of these solutions to fungi in terms of percentage inhibition of mycelial growth was calculated using the formula:

$$\% \text{ Inhibition} = (\text{dc} - \text{dt}) / \text{dc} \times 100$$

Where, dc = average increase in mycelial growth in control, dt = average increase in mycelial growth under treatment.

4. Results

4.1. FTIR Analysis

FTIR spectra of Thyme and Oregano EOs, Cs, CFs, CFs+EOs were recorded to investigate CFs and Thyme-Oregano EOs interactions. The results of FTIR spectra are shown in Figures 1 and 2, and the main bands are shown in Table 1. Thyme and oregano essential oils show characteristics peaks at 3397 and 3402 (hydrogen bonded OH stretching overlapped with N-H stretching bands), respectively. These peaks are joined, increased




Figure 1. FTIR spectra of (a) Oregano EO, (b) Thyme EO, (c) the blend of Thyme and Oregano EOs, and (d) EOs loaded-chitosan film. The peaks show the presence of interaction in between two EOs in mixture.

in the intensity of the OH and NH stretching in a single peak at 3443 indicating an increase in the content of ester groups when thyme and oregano EOs are mixed together (Fig. 1). The crosslink between EOs shows characteristics peaks at 1638 (amide I) and 723 (C-O stretching) resulting from many peaks found in the individual EO which merged to a single peak when EOs are mixed in tween 20. It should be noted that the OH and N-H stretching bands around 3443 cm^{-1} remained undisturbed when EOs are added in chitosan film solution.

Chitosan powder shows the characteristic peak at 3296 cm^{-1} corresponding to the combined peaks of the OH and NH₂ groups stretching vibration in chitosan

(Fig. 2). A change in the relative intensity of this band was observed when EOs was incorporated in the chitosan films solution. A shift from 3296 to 3443 cm^{-1} is shown in EOs-loaded chitosan film solution indicates that the hydroxyl groups and amine groups of the chitosan might involve in some interactions after incorporation of EOs to the chitosan. In chitosan powder, the bands at 1662 and 1596 cm^{-1} were attributed to amide I (C=O stretching) and amide II (N-H bending modes) groups, respectively, joined and shifted to 1636 in the EOs-loaded chitosan film solution suggesting the crosslink of EOs to nitrogen atoms of chitosan. Peaks found in chitosan powder at 895 and 664 cm^{-1} joined and shifted to a new band at 722 cm^{-1} in EOs-loaded chitosan film

Table 1. Main IR bands (Cm^{-1}) of Cs Powder, CFs+EOs, Thyme EO, Oregano EO, and a blend of Thyme and Oregano EOs.

Wave number (cm^{-1})	Thyme EO (v.cm^{-1})	Oregano EO (v.cm^{-1})	Thyme+Oregano EOs (v.cm^{-1})	Cs Powder (v.cm^{-1})	CFs+EOs (v.cm^{-1})
3000-3500	3397	3402	3443	3296	3443
2500-3000	2961-2926-	2961-2924	/	2873	/
1500-2000	1619-1584	1680-1618	1638	1662-1596	1636
1000-1500	1458-1380	1451-1374	/	1424-1382-	/
500-1000	738	808	723	895-664	722




Figure 2. FTIR spectra of (a) chitosan powder, (b) chitosan film, (c) chitosan film enriched with EOs, and (d) EOs. The peaks show that the hydroxyl groups and amino groups of the chitosan might be involved in some interactions after incorporation of EOs to the chitosan.

solution indicating the interaction between chitosan and EOs. The present results indicate that blend of thyme and oregano EOs might be incorporated into the chitosan film.

4.2. Antibacterial Activity

The combined EOs and chitosan solution exhibited a bactericidal effect against all bacterial strains with a variable degree of susceptibility. The MIC and MBC of EOs, CFs, and CFs enriched with EOs against *Escherichia coli*, *Staphylococcus aureus*, *Lactobacillus rahnmosus* and *Enterococcus faecium* are shown in Table 2. The MIC and MBC values for EOs, CFs, and CFs enriched with EOs against *Escherichia coli* and

Staphylococcus aureus is very low compared to the concentrations needed to inhibit the growth of beneficial bacteria, *Lactobacillus rahnmosus* and *Enterococcus faecium*. However, a combination of CFs enriched with EOs exhibited a strong antibacterial properties against *Escherichia coli* (MIC= 0.625 mg.mL⁻¹, MBC= 1.25 mg.mL⁻¹), while the inhibition was very low with CFs without EOs (MIC= 2.5 mg.mL⁻¹, MBC= 5 mg.mL⁻¹).

The useful Gram-positive bacteria, *Lactobacillus rahnmosus* and *Enterococcus faecium* were more resistant to EOs, CFs, and CFs enriched with EOs. The MIC of EOs and CFs were 25 µL.mL⁻¹ and 5 mg.mL⁻¹, while the MBC were more than 25 µL.mL⁻¹ and 10 mg.mL⁻¹, respectively. When the CFs was enriched


Table 2. MIC (µL.mL⁻¹) and MBC (µL.mL⁻¹) of the essential oils (EOs), chitosan film solution (CFs), and chitosan film solution enriched with EOs (CFs+EOs)^a.

Bacteria	EOs		CFs		CFs+Eos	
	MIC	MBC	MIC	MBC	MIC	MBC
<i>Escherichia coli</i>	0.39	0.78	2.5	5	0.625	1.25
<i>Staphylococcus aureus</i>	12.5	25	2.5	5	2.5	5
<i>Lactobacillus rahnmosus</i>	25	>25	5	>10	5	>10
<i>Enterococcus faecium</i>	25	>25	5	>10	2.5	5

Table 3. MIC ($\mu\text{L.mL}^{-1}$) and MFC ($\mu\text{L.mL}^{-1}$) of the essential oils (EOs), chitosan film solution (CFs), and chitosan film solution enriched with EOs (CFs+EOs)^a.

Fungi	EOs		CFs		CFs+EOs	
	MIC	MFC	MIC	MFC	MIC	MFC
<i>Aspergillus niger</i>	12.5	25	>10	>10	>10	>10
<i>Alternaria alternata</i>	12.5	25	>10	>10	5	10


with EOs, the MIC was reduced to 2.5 mg.mL^{-1} and the bactericidal concentration was 5 mg.mL^{-1} against *Enterococcus faecium*, while against *Lactobacillus rahnosus* the MIC was 5 mg.mL^{-1} and the bactericidal concentration was $>10 \text{ mg.mL}^{-1}$.

**Figure 3.** The percentage of microbial growth inhibition carried out against *Aspergillus niger* and *Alternaria alternata* by a blend of Thyme and Oregano essential oils.

4.3. Antifungal Activity

The results obtained on the effect of the blend of thyme and oregano EOs, CFs, and the combination of CFs with EOs has shown a significant inhibitory activity on the growth of *Aspergillus niger* and *Alternaria alternata*. These two fungi exhibited the same susceptibility to EOs with MIC of 12.5 mg.mL^{-1} and MFC of 25 mg.mL^{-1} (Table 3). In contrast, both *Aspergillus niger* and *Alternaria alternata* were found not to be sensitive to CFs at all concentrations (MIC and MFC $>10 \text{ mg.mL}^{-1}$). The percent inhibition of the two strains, *Aspergillus niger* and *Alternaria alternata* by EOs is shown in Figure 3. For all the fungal strains, the MFC that results into no fungal growth on the plates was found to be $25 \mu\text{L.mL}^{-1}$, while the growth inhibition was observed even with the smallest concentrations (0.39 and $0.195 \mu\text{L.mL}^{-1}$ respectively for *Aspergillus niger* and *Alternaria alternata*. *Alternaria alternata* seems to be more sensitive than *Aspergillus niger*.

Chitosan in solution without EOs does not show an inhibitory effect against *Aspergillus niger* at all concentrations (Fig. 4). Considering CFs enriched with the EOs, as the concentration was increased,

**Figure 4.** The percentage of microbial growth inhibition carried out against *Aspergillus niger* and *Alternaria alternata* by CFs and CFs+EOs solutions.

the antifungal effect was also exacerbated. At the concentration of $10 \mu\text{L}\cdot\text{mL}^{-1}$ of the EOs and $10 \text{mg}\cdot\text{mL}^{-1}$ of chitosan in the film solution, no fungal growth (100% inhibition) was noticed and the inhibition activity was observed against all the fungal strains even with the smallest concentrations of chitosan and EOs in the solution.

5. Discussion

When Thyme and Oregano EOs are mixed, FTIR spectra revealed characteristic peaks at 1638 (amide I) and 723 (C-O stretching) resulting from many peaks found in the individual EO which merged to a single peak. However, the OH and N-H stretching bands around 3443cm^{-1} remained undisturbed when the mixture of EOs are added to the chitosan film solution. This is consistent with the results of Hosseini *et al.* (16) that have recorded the same characteristic peaks in the FTIR spectra of the oregano EO at the same wave number in the EO-loaded chitosan nanoparticles indicating no modification or interaction between this EO and chitosan nanoparticles. In the present study, a shift from 3296 to 3443cm^{-1} is found in the EOs-loaded chitosan film solution indicating that the hydrogen bonding was affected by the binding of EOs to the chitosan. Similar results were reported by Regiel *et al.* (32) when chitosan in solution was chemically cross-linked with the silver. According to Hosseini *et al.* (16), the binding of the EO to chitosan in solution results in a marked increase in the intensity of the CH stretching peaks which indicates an increase in the content of ester groups coming from EO molecules.

It is well known that CFs exhibit antibacterial activity due to their well-developed surface, which provides maximum contact with the bacteria (31). The MIC and MBC values for EOs, CFs, and CFs enriched with EOs against *Escherichia coli* and *Staphylococcus aureus* is very low compared to the concentrations needed to inhibit the growth of useful bacteria such as *Lactobacillus rhamnosus* and *Enterococcus faecium*. In similar studies, Rodríguez-Núñez *et al.* (29) have reported a larger bactericidal effect of chitosan film solution against Gram-negative bacteria (*Salmonella typhimurium*) than Gram-positive (*Staphylococcus aureus*) bacteria; results that are consistent with those found in the present study. In this study, CFs enriched with EOs presented a strong antibacterial property against *Escherichia coli*, while the inhibition was very low with CFs alone (i.e., without EOs). Our observation is in agreement with the results obtained by other authors on the chitosan film with a similar degree of deacetylation; with preparation conditions either

similar or different (30-32).

This study has revealed that the useful Gram-positive bacteria, *Lactobacillus rhamnosus* and *Enterococcus faecium* were more resistant to EOs, CFs, and CFs enriched with EOs. It has been previously reported by López-Mata *et al.* (30) that chitosan films incorporated with carvacrol has high bactericidal property against the two major Gram-negative food contaminant pathogenic bacteria (i.e., *Escherichia coli* and *Salmonella typhimurium*). Carvacrol is a phenolic compound found in essential oils of oregano and thyme, possessing antimicrobial and antioxidants properties (1-2, 28). Chitosan has the ability to bind and improve the bioactive components and bactericidal activities of EOs (16). The high antibacterial activity that was recorded against pathogenic bacteria in the present study could be explained by the fact that the positively charged chitosan enriched with EOs can create a semi-permeable barrier capable of reducing respiration and microbial growth retardation (29-30, 33). This result can also be explained by the direct contact between the bacteria and the chitosan-EOs films which could establish the electrostatic bond between the negatively charged bacterial wall and cationic chitosan which has demonstrated a superior antibacterial action (33). The antibacterial mechanism of carvacrol has been studied already and it is known that such antibacterial property is due to the changes caused by the interaction with lipophilic components of the bacterial membrane. This interaction causes a change in the permeability of H^+ and K^+ , disrupts enzyme system, compromises the genetic material of the bacteria, form fatty acid hydroperoxide by the oxygenation of unsaturated fatty acid, damages the essential functions and causes bacteria death (33-34).

The present results have revealed a significant inhibitory activity of the blend of thyme and oregano EOs, CFs, and CFs enriched with EOs on the growth of *Aspergillus niger* and *Alternaria alternata*. These results are in line with those previously reported by Kaur *et al.* (25), who have shown that antifungal activity of the chitosan nanoparticles against *Aspergillus flavus*, *Alternaria alternata* and *Rhizoctonia solani* was enhanced when chitosan nanoparticles were loaded with metals like Cu and Zn compared to the nanoparticles of chitosan alone. The antifungal effect of CFs enriched with EOs was strengthened as the concentration was increased. This is consistent with the results of Prabu and Natarajan, (35) who have previously reported that the antifungal effect of chitosan isolated from *Padophthalmus vigil* was strengthened as the concentration was increased. Comparing these

results with the one obtained in the present study, it could be deduced that the value obtained for the fungal growth inhibition is slightly higher in our investigation compared to those of the previous reports (25, 35); differences probably being due to the higher deacetylation degree of the used chitosan and the presence of EOs in the CFs.

In conclusion, this study has evaluated the antimicrobial effect of the chitosan films solution enriched with EOs against two pathogenic bacteria (*Staphylococcus aureus* and *Escherichia coli*), two useful bacteria (*Enterococcus faecium* and *Lactobacillus rhamnosus*), as well as two molds fungi (*Aspergillus niger* and *Alternaria alternata*). It is interesting that we could find that pathogenic bacteria are more susceptible to antibacterial agents compared to the useful bacteria. It was also observed that, antimicrobial activities were enhanced when chitosan film solution was enriched through preparation of a blend of chitosan with thyme and oregano essential oils compared to the chitosan film solution alone. These results show that chitosan-EOs complexes could be the promising candidates as a potent antimicrobial inhibitory agent that could be used as a coating for the monogastric animal feeds.

Acknowledgements

We gratefully thank the research facility at NIGEB, Iran for antimicrobial and FTIR analysis. Mr Kana Jean Raphaël acknowledges the fellowship received from TWAS-UNESCO Associateship Scheme at Centres of Excellence in the South.

References

- Bento MHL, Ouwehand AC, Tiihonen K, Lahtinen S, Nurminen P, Saarinen MT, Schulze H, Mygind T, Fischer J. Essential oils and their use in animal feeds for monogastric animals - Effects on feed quality, gut microbiota, growth performance and food safety: a review. *Vet Med.* 2013; **58**(9): 449-458.
- Krishan G, Narang A. Use of essential oils in poultry nutrition: A new approach. *J Adv Vet Anim Res.* 2014; **1**(4): 156-162. DOI: 10.5455/javar.2014.a36
- Chao SC, Gary YD. Screening for inhibitory activity of essential oils on selected bacteria, fungi and viruses. *J Essent Oil Res.* 2000; **12**: 639-649. DOI: 10.1080/10412905.2000.9712177
- Cross DE, McDevitt RM, Hillman K, Acamovic T. The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in chickens from 7 to 28 days of age. *Brit Poult Sci.* 2007; **48**: 496-506. DOI: 10.1080/00071660701463221
- Kirsti T, Kettunen H, Bento MHL, Saarinen M, Lahtinen S, Ouwehand AC, Schulze H, Rautonen N. The effect of feeding essential oils on broiler performance and gut microbiota. *Brit Poult Sci.* 2010; **51**: 381-392. doi: 10.1080/00071668.2010.496446
- Cao PH, Li FD, Li YF, Ru YJ, Péron A, Schulze H, Bento H. Effect of essential oils and feed enzymes on performance and nutrient utilization in broilers fed a corn/soy-based diet. *Int J Poult Sci.* 2010; **9**(8): 749-755. doi: 10.3923/ijps.2010.749.755
- Khattak F, Ronchi A, Castelli P, Sparks N. Effects of natural blend of essential oil on growth performance, blood biochemistry, cecal morphology, and carcass quality of broiler chickens. *Poult Sci.* 2014; **93**: 132-137. DOI: 10.3382/ps.2013-03387
- Tiihonen K, Kettunen H, Bento MHL, Saarinen M, Lahtinen S, Ouwehand AC, Schulze H, Rautonen N. The effect of feeding essential oils on broiler performance and gut microbiota. *Brit Poult Sci.* 2010; **51**(3): 381-392. DOI: 10.1080/00071668.2010.496446
- Amerah AM, PE'Ron A, Zaefarian F, Ravindran V. Influence of whole wheat inclusion and a blend of essential oils on the performance, nutrient utilisation, digestive tract development and ileal microbiota profile of broiler chickens. *Brit Poult Sci.* 2011; **52**(1): 124-132. DOI: 10.1080/00071668.2010.548791
- Zhang KY, Yan F, Keen CA, Waldroup PW. Evaluation of microencapsulated essential oils and organic acids in diets for broiler chickens. *Int J Poult Sci.* 2005; **4**(9): 612-619. DOI:10.3923/ijps.2005.612.619
- Šegvić Klarić M, Kosalec I, Mastelić J, Piecková E, Pepeljnak S. Antifungal activity of thyme (*Thymus vulgaris* L.) essential oil and thymol against moulds from damp dwellings. *Lett Appl Microb.* 2007; **44**: 36-42. DOI: 10.1111/j.1472-765X.2006.02032.x
- Tullio V, Nostro A, Mandras N, Dugo P, Banche G, Cannatelli MA, Cuffini AM, Alonzo V, Carlone NA. Antifungal activity of essential oils against filamentous fungi determined by broth microdilution and vapour contact methods. *J Appl Microb.* 2007; **102**: 1544-1550. DOI: 10.1111/j.1365-2672.2006.03191.x
- Brenes A, Roura E. Essential oils in poultry nutrition: main effects and modes of action. *Anim Feed Sci Technol.* 2010; **158**: 1-14. DOI:10.1016/j.anifeedsci.2010.03.007
- Bassolé IHN, Juliani HR. Essential oils in combination and their antimicrobial properties. *Molecules* 2012; **17**: 3989-4006. DOI: 10.3390/molecules17043989
- Bustos R, Romo L, Yáñez K, Díaz G, Romo C. Oxidative stability of carotenoid pigments and polyunsaturated fatty acids in microparticulates diets containing krill oil for nutrition of marine fish larvae. *J Food Engin.* 2003; **56**: 289-293. DOI: 10.1016/S0260-8774(02)00272-8
- Hosseini SF, Zandi M, Rezaei M, Farahmandghavi F. Two-step method for encapsulation of oregano essential oil in chitosan nanoparticles: preparation, characterization and in vitro release study. *Carbohydr Poly.* 2013; **95**: 50-56. DOI: 10.1016/j.carbpol.2013.02.031
- Fernández-Urrusuno R, Calvo P, Remuñan-López C, Vila-Jato JL, Alonso MJ. Enhancement of nasal absorption of insulin using chitosan nanoparticles. *Pharm Res.* 1999; **16**(10):1576-81. DOI: 10.1023/A:1018908705446
- Vila A, Sánchez A, Janes K, Behrens L, Kissel T, Vila-Jato JL, Alonso MJ. Low molecular weight chitosan nanoparticles as new carriers for nasal vaccine delivery in mice. *Eur J Pharm Biopharm.* 2004; **57**(1):123-31. DOI: 10.1016/j.ejpb.2003.09.006
- Rezaei Mokarram A, Alonso MJ. Preparation and evaluation of chitosan nanoparticles containing diphtheria toxoid as new carriers for nasal vaccine delivery in mice. *Arch Razi Inst.* 2006; **61**(1):13-25.
- Mohammadpour Dounighi N, Behfar A, Ezabadi A, Zolfagharian H, Heydari M. Preparation of Chitosan nanoparticles containing

- Naja-naja oxiana* snake venom. *Nanomedicine*, 2010; **6**(1):137-43. DOI: 10.1016/j.nano.2009.06.002
21. Mohammadpour Dounighi N, Eskandari R, Avadi MR, Zolfagharian H, Mir Mohammad Sadeghi A, Rezayat M. Preparation and *in vitro* characterization of chitosan nanoparticles containing *Mesobuthus eupeus* scorpion venom as an antigen delivery system. *J Vernu Anitox Trop Dis*. 2012; **18**(1): 44-52. DOI: 10.1590/S1678-91992012000100006
 22. Qi L, Xu Z, Jiang X, Hu C, Zou X. Preparation and antibacterial activity of chitosan nanoparticles. *Carboh Res*. 2004; **339**: 2693-2700. doi:10.1016/j.carres.2004.09.007
 23. New N, Furuike T, Tamura H. The mechanical and biological properties of chitosan scaffolds for tissue regeneration templates are significantly enhanced by chitosan from *Gongronella butleri*. *Materials* 2009; **2**(2): 374-398. DOI:10.3390/ma2020374
 24. Wei D, Sun W, Qian W, Ye Y, Ma X. The synthesis of chitosan-based sliver nanoparticles and their antibacterial activity. *Carboh Res*. 2009; **344**: 2375-23. DOI: 10.1016/j.carres.2009.09.001
 25. Kaur P, Thakur R, Barnela M, Chopra M, Manuja A, Chaudhury A. Synthesis, characterization and *in vitro* evaluation of cytotoxicity and antimicrobial activity of chitosan-metal nanocomposites. *J Chem Technol Biotech*. 2015; **90**(5): 867-873. DOI: 10.1002/jctb.4383
 26. Vimal S, Abdul Majeed S, Taju G, Nambi KSN, Sundar Raj N, Madan N, Farook MA, Rajkumar T, Gopinath D, Sahul Hammed AS. Chitosan tripolyphosphate (CS/TPP) nanoparticles: Preparation, characterization and application for gene delivery in shrimp. *Act Trop*. 2013; **128**: 486-493. DOI: 10.1016/j.actatropica.2013.07.013
 27. Vimal S, Taju G, Nambi KSN, Abdul Majeed S, Sarath Babu V, Ravi M, Sahul Hammed AS. Synthesis and characterization of CS/TPP nanoparticles for oral delivery gene in fish. *Aquaculture*, 2012; **358/359**: 14-22. DOI:10.1016/j.aquaculture.2012.06.012
 28. Chi S, Zivanovic S, Penfield MP. Application of chitosan films enriched with oregano essential oil on Bologna-active compounds and sensory attributes. *Food Sci Technol Int*. 2006; **12**(2):111-117.
 29. Rodríguez-Núñez JR, López-Cervantes J, Sanchez-Machado DI, Ramirez-Wong B, Torres-Chavez P, Cortez-Rocha MO. Antimicrobial activity of chitosan-based films against *Salmonella typhimurium* and *Staphylococcus aureus*. *Int J Food Sci Tech*. 2012; **47**: 2127-2133. DOI: 10.1111/j.1365-2621.2012.03079.x. © 2012
 30. Lopez-Mata MA, Ruiz-Cruz S, Silva-Beltran NP, Ornelas-Paz JJ, Zamudio-Flores PB, Burruel-Ibarra SE. Physicochemical, antimicrobial and antioxidant properties of chitosan films incorporated with Carvacrol. *Molecules* 2013; **18**: 13735-13753. DOI: 10.3390/molecules181113735
 31. Leceta I, Guerrero P, Ibarburu I, Duenas MT, De la Caba K. Characterization and antimicrobial analysis of chitosan-based films. *J Food Eng*. 2013; **116**: 889-899. DOI: 10.1016/j.jfoodeng.2013.01.022
 32. Regiel A, Irusta S, Kyziol A, Arruebo M, Santamaria J. Preparation and characterization of chitosan-silver nanocomposite films and their antibacterial activity against *Staphylococcus aureus*. *Nanotechnology* 2013; **24**: 1-13. DOI: 10.1088/0957-4484/24/1/015101
 33. Ruiz-Navajas Y, Viuda-Martos M, Sendra E, Perez-Alvarez JA, Fernandez-Lopez J. *In vitro* antibacterial and antioxidant properties of chitosan edible films incorporated with *Thymus moroderi* or *Thymus piperella* essential oils. *Food Cont*. 2013; **30**: 386-392. DOI: 10.1016/j.foodcont.2012.07.052
 34. Ultee SA, Bennik M, Moezelaar R. The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Appl Environ Microb*. 2002; **68**: 1561-1568. DOI: 10.1128/AEM.68.4.1561-1568.2002
 35. Prabu K, Natarajan E. *In vitro* antimicrobial and antioxidant activity of chitosan isolated from *Podophthalmus vigil*. *J Appl Pharm Sci*. 2012; **2**(9): 75-82. DOI: 10.7324/JAPS.2012.2916