

## Brief Report



# Differential Expression of Mitochondrial *Manganese Superoxide Dismutase (SOD)* in *Triticum aestivum* Exposed to Silver Nitrate and Silver Nanoparticles

Javad Karimi <sup>1,2\*</sup>, Sasan Mohsenzadeh <sup>2</sup>, Ali Niazi <sup>3</sup> and Ali Moghadam <sup>3</sup>

<sup>1</sup> Department of Biotechnology, Faculty of Biological Sciences and Technology, Shahid Ashrafi Esfahani University, Isfahan, Iran

<sup>2</sup> Department of Biology, Faculty of Sciences, Shiraz University, Shiraz 71454, Iran

<sup>3</sup> Institute of Biotechnology, Shiraz University, 71441-65186 Bajgah, Shiraz, Iran

\*Corresponding author: Javad Karimi, Department of Biotechnology, Faculty of Biological Sciences and Technology, Shahid Ashrafi Esfahani University, Isfahan, Iran and Department of Biology, Faculty of Sciences, Shiraz University, Shiraz 71454, Iran.  
Tel: +98 71 36137737, E-mail: javadkarimi@shirazu.ac.ir

Received: 9 September 2015; Revised: 4 October 2017; Accepted: 7 October 2017; Published online: 29 December 2017

**Background:** The increasing use of nanoparticles (NPs) may have negative impacts on both organisms and the environment.

**Objectives:** The differential expression of mitochondrial manganese superoxide dismutase (*MnSOD*) gene in wheat in response to silver nitrate nanoparticles (AgNPs) and AgNO<sub>3</sub> was investigated.

**Materials and Methods:** A quantitative Real-Time RT-PCR experiment was carried out with *MnSOD* gene using RNAs isolated from wheat shoots treated for 0, 2, 6, 12, and 24 h with 100 mg.L<sup>-1</sup> of either AgNO<sub>3</sub> or AgNPs.

**Results:** The results of this study showed that both treatments cause changes in the expression pattern of the *MnSOD* gene. While 2 and 6 h following the beginning of the stress, *MnSOD* expression was up-regulated significantly, in response to AgNO<sub>3</sub> (1.4 and 2.8 fold, respectively), in response to AgNPs, it was up-regulated significant only after 6 h (1.6 fold), compared with the control. The gene expression, after 12 h in response to AgNO<sub>3</sub> and AgNPs were down-regulated significantly (0.7 and 0.8 fold, respectively), and in the next 12 h, the expression appeared to be similar to the control.

**Conclusion:** Exposure to both AgNPs and Ag ions led to a significant increase in *MnSOD* expression, but AgNO<sub>3</sub> changed the *MnSOD* expression faster than AgNPs. Therefore, it is suggested that AgNO<sub>3</sub> has greater penetrability and effectiveness.

**Keywords:** Gene expression, Silver nanoparticles, Silver nitrate, Superoxide dismutase, *Triticum aestivum*.

## 1. Background

Today, nanoparticles, because of their unusual, novel, and improved properties, are applied in all aspects of life. In recent years, design, production, optimization, and application of synthesized nanoparticles in different areas and mode of functions are very attractive (1, 2). The growing production of nanoparticles has led to concerns over the adverse impacts on the environment (3, 4). Based on literature there is no full understanding of the potential for health or environmental risks on nanoparticles (5). The effects of metal nanoparticles

on plant germination and growth with focus on crop yield have been reported elsewhere (6-10). In contrast, possible side effects of the use of silver nanoparticles and silver ions on plant growth has received little attention (11-14).

Both silver nanoparticles (AgNPs) and ions (AgNO<sub>3</sub>) have been negative cellular and molecular effects on plants such as interaction with the cell wall and cell membrane, leading to changes in membrane permeability, impairing proton motive force and inhibition of the ATP synthesis, interactions with amino

acids and inhibition of enzyme activity by binding to sulfhydryl groups (-SH) of amino acids and their active center; inhibition of electron movement in the respiratory chain and cytochromes; deregulation of DNA and RNA synthesis; denaturing the ribosome and inhibiting protein synthesis; and generation of reactive oxygen species (ROS) (8, 15-17).

In plants, reactive oxygen species (ROS) are produced and scavenged in both stressed and unstressed conditions. In non-stressed condition the production and recycling the ROS appears to be in balance, while in stressed condition the production supersedes the scavenging and therefore the cascades of cellular and molecular damages will be rolled over throughout the organism (18). Under these stress, with high concentration of ROS, an effective scavenging system is required to avoid oxidative damage to plants (18). *Superoxide dismutase (SOD)* enzyme has been reported to serve as an environmental toxicological stress marker, which plays a determinant role in plant protection against the toxic effects of oxidative stress by scavenging superoxide radicals. *SOD* catalyzes the conversion of  $O_2^-$  into  $H_2O_2$ , which can reduce the amounts of damaging ROS produced by the adverse environments, and alleviate the harm to plants (19-21). In plants, multiple *SOD* isozymes exist, which are classified into three types based on their metal cofactors: *Cu/ZnSOD*, *FeSOD* and *MnSOD* (18). To the best of our knowledge, limited investigations have been conducted to study the effects of NPs on plant gene expression. The release of NPs in the environment has increased concerns about their effects on living organisms, including higher plants (22, 23).

## 2. Objectives

Here, the changes of *MnSOD* were studied in wheat exposed to both AgNPs and  $AgNO_3$  via Real-Time PCR.

## 3. Materials and Methods

### 3.1. Plant Materials

Wheat seeds (*Triticum aestivum* L. Var. Chamran) were obtained from Zarghan Agricultural Research Center, Iran and kept in the dark at 4 °C before use. Seeds were surface sterilized by soaking in 5% (w/v) sodium hypochlorite for 10 min. They were washed three times with distilled water and air-dried on filter papers. Seeds were allowed to germinate in dark at 25 °C on moist filter papers. Five-day old seedlings (20) were transferred into small plastic containers filled with perlite and Hoagland nutrient solution (pH 6.2). Wheat seedlings

were grown in growth chamber set at 16 h/8 h light-dark periods. Three biological and technical replicates were used. Control and treated wheat seedlings were immediately frozen in liquid nitrogen and stored at -80 °C until nucleic acid extraction.

### 3.2. Silver Nitrate and Silver Nanoparticles Treatments

AgNPs with average sizes of 20 nm and 99.99% purity was purchased from US Research Nanomaterials, Inc. (USA).  $AgNO_3$  (99.9%) was obtained from Sigma-Aldrich (Sigma-Aldrich-209139, Germany). Using Hoagland nutrient solution as solvent, two concentrations (10 and 100  $mg.L^{-1}$ ) of both silver particles were prepared. The dissolved particles were dispersed by a high-power probe-type Sonicator (Misonix, Qsonica LLC, Newton, USA) for 30 min. For the control only the Hoagland nutrient solution was used. After 2, 6, 12 and 24 h of treatment, the leaves of wheat were harvested and were stored at -80 °C until RNA extraction.

### 3.3. RNA Isolation and cDNA Synthesis

Total RNA was extracted using an RNA isolation kit (DNA Zist kit, Iran) according to the manufacturer's instructions. Extracted RNA was kept in -80 °C until DNaseI treatment (Fermentas, Hanover, MD). Extracted RNA was quantified using Nano-Drop ND 1000 Spectrophotometer (Wilmington, USA). *Actin* (accession numbers: GQ339780 and KC775782) and *18s rRNA* (accession number: K01229) primers (Table 1) were used to check for any DNA contamination. First strand cDNA was synthesized using reverse transcriptase (Fermentas, Hanover, MD).

### 3.4. Primer Design

*MnSOD* primers (Table 1) were designed at Allele ID 7.8 software according to the relevant wheat sequences (accession numbers: KC158224, AY963808, KP313757) to amplify 105 fragment. The wheat *Actin* and *18s rRNA* genes were used as the reference genes (24) for data normalization via geometric mean. PCR products after preparation were sequenced in Macrogen. Sequence homology searches were carried out using the Basic Local Alignment Search Tool (BLAST) facility available at NCBI.

### 3.5. Real-Time RT-PCR

Real-Time RT-PCR reactions were performed using a SYBR Green kit (Takara, Japan) according to the manufacturer's protocol. The first strand cDNA were diluted 5X with RNase-free water. Quantitative PCR was performed in 20  $\mu L$  reactions using gene specific

**Table 1.** Primers used for Real-Time RT-PCR amplification, Amplicon size and T<sub>m</sub>. The accession numbers of wheat *Actin*, were GQ339780 and KC775782 and accession number of *18s rRNA* gene was K01229.

Genes	Primer Sequence 5'→3'	Product length (bp)	T <sub>m</sub> (°C)
<i>SOD</i>	F: TCCGCCGTCGTCCACCTC	105	63.8
	R: CCACCACCTCGCTGATG		
<i>Actin</i>	F: GACATACAATTCCATCAT	153	56
	R: TTAACCTTCATACTGCTA		
<i>18s rRNA</i>	F: CGCTCCTACCGATTGAATGG	127	56
	R: CCTTGTTACGACTTCTGCTTCC		

primers (A final primer concentration of 0.4 μM is recommended in most reactions), 4 μL of cDNA as a template. The abundance of targeted gene transcripts was normalized to Actin and 18s rRNA and set relative to control plants according to the  $2^{-\Delta\Delta CT}$  method (25). For quantitative Real-Time RT-PCR data, relative expressions were calculated based on the threshold cycle (CT) method.

### 3.6. Statistical Analysis

Experiment was conducted in the frame of randomized complete block design and each value reported was the average of three replicates. The raw data were imported into Microsoft Excel 2007 and GraphPad Prism 5 programs for calculations and graphic representation. SPSS (version 16.0) software was used for the analysis of variance. Quantitative changes of parameters were evaluated through analysis of variance (two-way ANOVA), with Duncan's multiple range tests at  $p \leq 0.05$  to find out significant differences among treatments. All results are presented as the means  $\pm$  standard deviation (SD).

## 4. Results

### *Analysis of MnSOD Expression*

Real-Time RT-PCR was performed to investigate the *MnSOD* expression in response to AgNPs and AgNO<sub>3</sub>. The results showed that the *MnSOD* was significantly up-regulated at 2 and 6 h (1.3 and 2.8 fold change, respectively) and then down-regulated at 12 h (0.5 fold) after AgNO<sub>3</sub> treatment (Table 2 and Fig. 1). Twenty four hours after AgNO<sub>3</sub> treatment, the expression of *MnSOD* remained unchanged. In response to AgNPs, *MnSOD* expression increased at 6 h (1.6 fold) and down-regulated at 12 h (0.8 fold) compared with non-treated wheat seedling at the same times. At 2 and 24 h after AgNPs treatment, there were no significant changes


as compared to the control. *MnSOD* expression was gradually increased at 2 h after treatment, and reached to its maximum level at 6 h post treatment, in response to both AgNO<sub>3</sub> and AgNPs. The *MnSOD* expression was decreased slightly at 12 h of post-treatment and reached to the control level, during next 12 h. Results showed that *MnSOD* gets up-regulated early in response to silver in two forms of ions and nanoparticles.

## 5. Discussion

It is not quite clear, where the toxicity, comes from, during the release of silver nanoparticles; Is it from the nano-molecules or from the release of silver itself? In this study, the expression of *MnSOD* was surveyed in *T. aestivum* seedlings exposed to silver ions and silver nanoparticles. Here, we observed that exposure to both AgNPs and Ag ions led to a significant increase in

**Table 2.** The relative expression of *MnSOD* in wheat seedlings exposed to AgNPs and AgNO<sub>3</sub> (control, and 100 mg.L<sup>-1</sup>) at different time intervals post-treatment (0, 2, 6, 12, and 24 h).

Treatment (mg.L <sup>-1</sup> )	SOD expression
AgNO <sub>3</sub> 2 h	1.44 $\pm$ 0.10 <sup>b</sup>
AgNPs 2 h	1.05 $\pm$ 0.10 <sup>c</sup>
Control 6 h	0.89 $\pm$ 0.07 <sup>cd</sup>
AgNO <sub>3</sub> 6 h	2.56 $\pm$ 0.12 <sup>a</sup>
AgNPs 6 h	1.47 $\pm$ 0.11 <sup>b</sup>
Control 12 h	0.86 $\pm$ 0.06 <sup>d</sup>
AgNO <sub>3</sub> 12 h	0.51 $\pm$ 0.04 <sup>f</sup>
AgNPs 12 h	0.69 $\pm$ 0.07 <sup>e</sup>
Control 24 h	0.78 $\pm$ 0.07 <sup>de</sup>
AgNO <sub>3</sub> 24 h	0.83 $\pm$ 0.05 <sup>de</sup>
AgNPs 24 h	0.75 $\pm$ 0.07 <sup>de</sup>



**Figure 1.** Analyses of the relative gene expression of *MnSOD* in wheat seedlings exposed to AgNPs and AgNO<sub>3</sub> (control, and 100 mg.L<sup>-1</sup>) and different times after treatment (0, 2, 6, 12, and 24 h). All results are presented as the means ± standard deviation (SD) and different letters above the bars indicate significant differences between treatments ( $p \leq 0.05$ ).

*MnSOD* expression. This gene was induced significantly and up-regulated by oxidative stress of both heavy metals, presumably applied in two different forms. Although AgNPs increased the expression of *MnSOD*, its effect was weaker than that of silver ion, consistent with earlier reports (11-14). In response to AgNO<sub>3</sub>, at 2 h of post-treatment, expression of *MnSOD* was up-regulated. Whereas in response to silver nanoparticle it was up-regulated 6 h after treatment. These data provided further evidences that the effects of AgNPs are not simply due to the release of Ag ions. These studies have led to new insights into the changes of *MnSOD* expression of wheat responses to AgNPs and AgNO<sub>3</sub>. Plants have an antioxidant protection system to keep away from ROS formation, and this protection system includes a range of antioxidant enzymes such as *SOD*; something that we can implement as the biomarker of the oxidative stress to analyze cellular homeostasis (26, 27). Our results suggest that wheat seedlings might develop strategies to cope with AgNO<sub>3</sub> and AgNPs toxicity by manipulating the expression level of *SOD* genes.

## 6. Conclusion

Exposure to both AgNPs and Ag ions led to a significant increase in *MnSOD* expression, but AgNO<sub>3</sub> changed the *MnSOD* expression faster than AgNPs. Therefore, it is suggested that AgNO<sub>3</sub> has greater penetrability and effectiveness. These findings suggest that the *MnSOD* expression can be considered as a marker for evaluation of the heavy metals and metal nanoparticles effects on the cellular processes. To better understand of AgNO<sub>3</sub> and AgNPs toxicity on plant, additional

researches in other genes that are important in this regard, should be carried out.

## Acknowledgment

The authors thank the Biology Department and Institute of Biotechnology of the Shiraz University for Supporting of this work.

## Conflicts of Interest

The authors declare no conflict of interest.

## References

- Albanese A, Tang PS, Chan WC. The effect of nanoparticle size, shape, and surface chemistry on biological systems. *Annu Rev Biomed Eng.* 2012;**14**:1-16. DOI: 10.1146/annurev-bioeng-071811-150124
- Gao J, Gu H, Xu B. Multifunctional magnetic nanoparticles: design, synthesis, and biomedical applications. *Acc Chem Res.* 2009;**42**(8):1097-107. DOI: 10.1002/adma.201000260
- Dreher KL. Health and environmental impact of nanotechnology: toxicological assessment of manufactured nanoparticles. *Toxicol Sci.* 2004;**77**(1):3-5. DOI: 10.1093/toxsci/kfh041
- Neal AL. What can be inferred from bacterium–nanoparticle interactions about the potential consequences of environmental exposure to nanoparticles? *Ecotoxicology.* 2008;**17**(5):362-71. DOI: 10.1007/s10646-008-0217-x
- Morgan K. Development of a preliminary framework for informing the risk analysis and risk management of nanoparticles. *Risk Anal.* 2005;**25**(6):1621-35. DOI: 10.1111/j.1539-6924.2005.00681.x
- Joblonski D. Evolutionary consequences of mass extinctions. In: Raup DM, editor. *Pattern and Process in the History of Life.* 1986;382-418. DOI: 10.1007/978-3-642-70831-2
- Khot LR, Sankaran S, Maja JM, Ehsani R, Schuster EW. Applications of nanomaterials in agricultural production and crop protection: a review. *Crop Prot.* 2012;**35**:64-70. DOI:



- 10.1016/j.cropro.2012.01.007
8. Nair R, Varghese SH, Nair BG, Maekawa T, Yoshida Y, Kumar DS. Nanoparticulate material delivery to plants. *Plant Sci.* 2010;**179**(3):154-63. DOI: 10.1016/j.plantsci.2010.04.012
  9. Rai M, Ingle A. Role of nanotechnology in agriculture with special reference to management of insect pests. *Appl Microbiol. Biotechnol.* 2012;**94**(2):287-93. DOI: 10.1007/s00253-012-3969-4
  10. Anderson A, McLean JE, Jacobson AR, Britt D. CuO and ZnO Nanoparticles Modify Interkingdom Cell Signaling Processes Relevant to Crop Production: *J Agri Food Chem.* 2017; In Press. DOI: 10.1021/acs.jafc.7b01302
  11. Qian H, Peng X, Han X, Ren J, Sun L, Fu Z. Comparison of the toxicity of silver nanoparticles and silver ions on the growth of terrestrial plant model *Arabidopsis thaliana*. *J Environ Sci.* 2013;**25**(9):1947-56. DOI: 10.1016/S1001-0742(12)60301-5
  12. Kaveh R, Li Y-S, Ranjbar S, Tehrani R, Brueck CL, Van Aken B. Changes in *Arabidopsis thaliana* gene expression in response to silver nanoparticles and silver ions. *Environ Sci Technol.* 2013;**47**(18):10637-44. DOI: 10.1021/es402209w
  13. Jiravova J, Tomankova KB, Harvanova M, Malina L, Malohlava J, Luhova L, *et al.* The effect of silver nanoparticles and silver ions on mammalian and plant cells in vitro. *Food Chem Toxicol.* 2016;**96**:50-61. DOI: 10.1016/j.fct.2016.07.015
  14. Tomacheski D, Pittol M, Simoes D, Ribeiro V, Santana R. Impact of silver ions and silver nanoparticles on the plant growth and soil microorganisms. *Global J Environ Sci Manage.* 2017;**3**(4):341-350. DOI: 10.22034/gjesm.2017.03.04.00\*
  15. Kaegi R, Voegelin A, Sinnet B, Zuleeg S, Hagedorfer H, Burkhardt M, *et al.* Behavior of metallic silver nanoparticles in a pilot wastewater treatment plant. *Environ Sci Technol.* 2011;**45**(9):3902-8. DOI: 10.1021/es10418922019.
  16. Kumari M, Mukherjee A, Chandrasekaran N. Genotoxicity of silver nanoparticles in *Allium cepa*. *Sci Total Environ.* 2009;**407**(19):5243-6. DOI: 10.1016/j.scitotenv.2009.06.024
  17. Monica RC, Cremonini R. Nanoparticles and higher plants. *Caryologia.* 2009;**62**(2):161-5. DOI: 10.1080/00087114.2004.10589681
  18. Alscher RG, Erturk N, Heath LS. Role of *superoxide dismutases (SODs)* in controlling oxidative stress in plants. *J Exp Bot.* 2002;**53**(372):1331-41. DOI: 10.1093/jxbbot/53.372.1331
  19. Ben-qin W. The Research Progress of the Plant *SOD*. *J Hebei Agr Sci.* 2008;**21**(4):6-9. DOI: 10.3969/j.issn.1088-1631.2008.03.003
  20. Bowler C, Montagu Mv, Inze D. Superoxide dismutase and stress tolerance. *Annu Rev Plant Biol.* 1992;**43**(1):83-116. DOI: 10.1146/annurev.pp.43.060192.000503
  21. Verma S, Dubey R. Lead toxicity induces lipid peroxidation and alters the activities of antioxidant enzymes in growing rice plants. *Plant Sci.* 2003;**164**(4):645-55. DOI: 10.1016/S0168-9452(03)00022-0
  22. Liu J, Hurt RH. Ion release kinetics and particle persistence in aqueous nano-silver colloids. *Environ Sci Technol.* 2010;**44**(6):2169-75. DOI: 10.1021/es9035557
  23. Sutherland WJ, Clout M, Cote IM, Daszak P, Depledge MH, Fellman L, *et al.* A horizon scan of global conservation issues for 2010. *Trends Ecol. Evol.* 2010;**25**(1):1-7. DOI: 10.1016/j.tree.2009.10.003
  24. Nicot N, Hausman J-F, Hoffmann L, Evers D. Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. *J Exp Bot.* 2005;**56**(421):2907-14. DOI: 10.1093/jxb/eri285
  25. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the Method. *Methods.* 2001;**25**(4):402-8.
  26. Gill SS, Tuteja N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol Biochem.* 2010;**48**(12):909-30. DOI: 10.1016/j.plaphy.2010.08.016
  27. Das K, Roychoudhury A. Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. Redox Homeostasis Managers in Plants under Environmental Stresses. *Front. Environ. Sci.* 2014;**2**:53-67. DOI: 10.3389/fenvs.2014.00053