

Functional analysis of Glycin Rich- RNA Binding protein, a suppressor of Trehalose-6-Phosphate mediating growth arrest in *Arabidopsis thaliana*

Mahnaz Aghdasi^{1,2*}, Henriette Schluempmann²

¹Department of Biology, Faculty of Science, Golestan University, P.O.Box 155, Gorgan, I.R. Iran ²Department of Molecular Plant Biology, Institute of Environmental Biology, Utrecht University, Padualaan 8, 3584 CH, The Netherlands

Abstract

Metabolism of the alpha-1,1 glucose disaccharide, trehalose, is indispensable in plants. In the Murashige and Skoog (MS) medium, trehalose inhibits plant growth and allocation of carbon to roots. A suppressor of trehalose-6-phosphate (T6P) mediated growth arrest, *GR-RBP2*, is characterized in more detail. Phylogenetic analysis revealed that *GR-RBP2* is a protein of likely prokaryotic origin. A knockout mutant of *GR-RBP2* was identified in the T-DNA insertion line SALK-059714, yet plants of this line were not altered with regard to growth on different carbon sources and on trehalose compared to WT. GUS expression analysis showed that *GR-RBP2* was detected in adult leaves, flowers and siliques. Expression was particularly high in root tips. *GR-RBP2* expression also is insensitive to 100 mM trehalose. TAP-tagged versions of this protein showed that *GR-RBP2* is part of a protein complex in *planta*.

Keywords: *Arabidopsis*; *GR-RBP2*; Gus fusion; Tap-tagged; Trehalose

INTRODUCTION

Trehalose is the alpha, alpha-1,1-linked glucose disaccharide, which is found in a wide variety of organisms (Elbein *et al.*, 2003). Its metabolism has recently been recognized to play an important role in carbon signaling in plants (Paul *et al.*, 2008; Rolland *et al.*, 2006).

Genes coding for the components involving in the trehalose metabolism have been identified in *Planta* recently. Synthesis of trehalose in plants is typically via its phosphorylated intermediate, trehalose-6-phosphate (T6P). Trehalose-6-phosphate synthase (TPS) converts UDP-glucose and glucose-6-phosphate to T6P. Trehalose phosphate phosphatase (TPP) dephosphorylates T6P to trehalose. Trehalase cleaves trehalose to two glucose molecules. There are 11 TPS and 10 TPP homologues in *Arabidopsis*. Whilst only one gene encodes trehalase (Vandesteene *et al.*, 2010; Pramanic and Imai 2005; Leyman *et al.*, 2001). Minor alterations of T6P steady states in plants, yield dramatic and pleiotropic phenotypic changes (Pramanic and Imai 2005; Pellny *et al.*, 2004; Schluempmann *et al.*, 2003). Additionally, deletion of the T6P synthase (TPS), *AtTPS1*, in *Arabidopsis* is lethal and can be overcome by complementation with active TPS enzyme (Schluempmann *et al.*, 2003; Eastmond *et al.*, 2002). Current evidences suggest an important regulatory role for T6P such as carbon utilization in *Arabidopsis* seedlings (Schluempmann *et al.*, 2004). It is not understood however, how T6P controls carbon utilization and why changes in the steady state level of T6P yield such strong phenotypic changes.

Trehalose feeding to *Arabidopsis* seedlings inhibits root growth, emergence of priming leaves and allocation of carbon to the root and shoots by a negative feedback (Aghdasi and Schluempmann 2009; Ramon *et al.*, 2007; Fritzius *et al.*, 2001; Wingler *et al.*, 2000). Increased trehalose and T6P also affect starch synthesis via the control of T6P over AGPase

*Correspondence to: Mahnaz Aghdasi, Ph.D.
Tel: +98 171 2327520; Fax: +98 171 2322906
E-mail: Aghdasi46@yahoo.com; M.aghdasi@gu.ac.ir

redox activation (Kolbe *et al.*, 2005). Trehalose feeding induces massive starch accumulation in sources tissues and inhibits starch accumulation in the columella cells of root tips (Aghdasi and Schluempmann 2009; Ramon *et al.*, 2007; Wingler *et al.*, 2000).

To understand how trehalose-6-phosphate control over growth, mutants were identified that accumulate similar amounts of T6P as WT, but that grows on trehalose medium, unlike WT. The screening has been performed in the Leclere and Bartel mutant collection (Leclere and Bartel, 2001). Interestingly, two of the trehalose resistant (*trr*) mutants express glycin-rich RNA-binding protein2 (GR-RBP2). Results from expression analysis and re-transformation with the CaMV35S/cDNA constructs showed that over-expression of *GR-RBP2* suppress trehalose-6-phosphate mediated growth arrest (Aghdasi *et al.*, 2007).

Proteome analysis revealed that GR-RBP2 is localized into the mitochondria of *Arabidopsis* (Kruft *et al.*, 2001). The GR-RBP2 protein is only 153 amino acids long. GR-RBP2 has a mitochondrial targeting sequence (amino acid residues 1-33). Amino acid residues 34-113 contain the RNA recognition motif (RRM) domain, with the consensus sequences ribonucleoprotein (RNP)-1 and RNP-2 RNA binding domains (Nomata *et al.*, 2004). Residues 122-153 contain the glycin-rich domain thought to be involved in protein-protein interactions (Lorkovic and Barta, 2002). The typical RRM consists of four anti-parallel β -strands and two α -helices arranged in a β - β , α - β fold with side chains that stack with RNA bases. RRM proteins are essential for regulation of post-transcriptional processes. The *Arabidopsis* genome encodes 200 RRM-containing proteins, among which eight proteins are classified into GR-RBP1-8 (Lorkovic, 2009; Lorkovic and Barta, 2002). Expression of several of this GR-RBP is induced by various stresses. Meanwhile it has been shown that GR-RBP3 has non-specific RNA and DNA binding ability (Carpenter *et al.*, 1994; Fang *et al.*, 1991; Condit and Meagher, 1987).

By using a reverse genetic approach, we attempted to identify processes that are controlled by T6P in plants. Phylogenetic analyses were carried out to identify GRRBP2 homologues in *Arabidopsis*. In an attempt to reveal GR-RBP2 protein function, identified T-DNA insertion lines were characterized. Transcriptional fusion constructs were made to visualize expression pattern of GR-RBP2. Tagged versions of GR-RBP2 were expressed to identify potential protein complexes with these proteins.

MATERIALS AND METHODS

Phylogenetic analyses: Protein sequences of GR-RBP2 were obtained from the TAIR database (WWW.Arabidopsis.org). The BLAST searches were performed, using the TAIR and NCBI BLAST programs at [http:// Arabidopsis.org/](http://Arabidopsis.org/) and <http://ncbi.gov/>, respectively. The multiple sequence alignment and the respective phylogenetic tree were performed at <http://www.ebi.ac.uk/clustalw/>, using the ClustalW method, scoring matrix Blosum, gap penalty of 10.

Plant materials: Seeds obtained from *Arabidopsis* transformed plants were sterilized by the gas-sterilization method (Salinas-Jose and Sanchez-Serrato, 2006). *Arabidopsis* seedlings were grown in agarose solidified (0.8%) half strength Murashige and Skoog (Murashige and Skoog, 1962) medium supplemented with antibiotics as required for selection, 100 mg/l gentamycin, 100 mg/l spectinomycin, 12.5 μ g/l hygromycin and 50 mg/l kanamycin. Plants were grown under long-day conditions at 22°C in soil.

Vector construction and plant transformation: To detect the expression pattern of *GR-RBP2*, fragments containing 2 kb promoters and the 5' UTRs of this gene was fused to the GUS sequence in vector pMDC162 (Curtis and Grossniklaus, 2003). The genomic sequences were PCR amplified from Col-0 using a two step procedure. In step one, amplification of template DNA was done by using template specific primers (Forward: 5'-AAAAAGCAGGCTTCGGTCTACTTACCTCCAGAT-3' and Reverse: 5'-AGAAAGCTGGGTATCTACAAAGTTCCAAAACAC-3'). Amplification conditions were performed as initial DNA denaturation at 95°C for 2 minutes followed by 10 cycles of 15 second denaturation at 94°C, 30 second annealing at 56°C and 2 minutes of extension at 72°C with a final extension time at 72°C for 10 minutes. Then 10 μ l of the first PCR product was used in the second PCR step, as a template. In this step, attB1 and attB2 adapter primers (Forward: 5'-GGGGA-CAAGTTTGTACAAAAAGCAC-3' and Reverse: 5'-AGGGACCACTTTGTACAAGAAAGCTGGGT-3') were used. The PCR amplification is carried out in two phases. The initial denaturing was done at 95°C for 2 min, followed by 5 cycles of 94°C for 15 second, annealing at 45°C for 30 second and extension at 72°C for 2 minute. The second amplification was done at 94°C for 15 second, annealing at 55°C for 30 second and extension at 72°C for 2 minutes and a final extension at 72°C for 10 minutes. PCR product was separat-

ed on agarose gel and the specific band was cut and cleaned using a DNA purification kit (Amersham, Biosciences, England).

PCR amplified fragments were cloned into pDONR201 by using the GateWay^R cloning system and the sequence was confirmed by DNA sequencing. Transferring fragments from pDONR201 to pMDC162 (*GUS* reporter gene) vector was performed using the recombination reaction as described by the manufacturer (Invitrogen, Carlsbad, CA). The construct was transformed into WT Col-0 plants by the floral dip method (Clough and Bent 1998) using *Agrobacterium tumefaciens* pGV2260. Transformed seedlings were selected on ½ MS supplemented with 12.5 µg/l hygromycin. The transformed seedlings were further grown on soil in long day conditions (16 h light, 8 h dark) at 22°C to obtain seed.

Translational fusions with the Tandem Affinity Purification (TAPa)-tag: To generate C-terminal fusions with the TAPa-tag, the cDNA's encoding the full-length versions of GR-RBP2 were amplified by PCR and cloned into pDONR201 (Rubio *et al.*, 2005). The transfer of genes from pDONR201 to TAPa vector was performed using the recombination reaction as described by the manufacturer (GateWay, Invitrogen, Carlsbad, USA). Constructs were transferred to wild-type plants (Col-0) by the floral dip method (Clough and Bent 1998). Transgenic seedlings were selected on solidified ½ MS medium supplemented with 100 mg/l gentamycin. The selected lines were grown on soil to obtain seed.

Western blot analyses: Protein fractions separated by SDS-PAGE (12% polyacrylamide) were blotted to polyvinylidene membrane (Millipore, Bedford, USA) in running buffer (3g Tris, 14.2 g glycine, 200 ml EtOH, 0.8 g SDS per liter) for 1.5 hour at 100 V, on ice. The membranes were blocked in phosphate buffered saline (PBS) supplemented with 5% w/v Milk (PBSM), incubated overnight with a 3000 fold dilution of Anti-myc primary antibody from mouse (Santa Cruz, biotechnology, Santa Cruz, CA) in PBSM. Unbound antibody was washed 4 times with PBS supplemented with 0.1% v/v Tween 20 (Merk, Hohenbrunn, Germany). The blots were incubated 1.5 hours with a 3000 fold dilution of anti-mouse IgG secondary antibody conjugated to horseradish peroxidase in PBSM. Unbound antibody was washed 4 times in PBS with 0.1% v/v Tween 20, before a one minute reaction with enhanced chemi-luminescence reagent (Sigma, USA) at room temperature and exposure of

the blots to X-ray film for 5 minutes (Kodak, Sigma-Aldrich, Germany).

Gel filtration on Sepharose S-300: Protein was extracted from frozen ground leaf material (0.5g) in 500 µl ice-cold buffer containing 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10% v/v glycerol, 0.1% w/v Nonidet P-40, 1 mM DTT, 1 mM PMSF, and 1× complete protease inhibitor cocktail (Roche, Indianapolis, IN, USA). The resulting crude extract was filtered through four layers of cheesecloth, and then centrifuged at 12,000 ×g (Eppendorf, UK) for 10 min at 4°C. The supernatant fraction (300 µg) was loaded onto a Superdex S-300 column (Amersham Biosciences, England) equilibrated in PBS and a flow rate of 0.1 ml/min. In each 1 ml, sixty fractions were collected. Proteins in each 3rd 1ml fraction were concentrated using 10 ul Strataresin (Stratagene company, La Jolla, USA). Resin bound proteins (5 ul) was fractionated by SDS-PAGE using 12% w/v polyacrylamide gels, that were either stained with Coomassie blue (R250, Sigma) or blotted for detection of the TAP-tag as described in the previous section.

Tandem affinity purification: TAPa-tagged proteins were purified essentially as described by Rubio *et al.* (2005). For protein extraction, fifteen gram of fresh leaves was harvested, snap frozen ground in liquid nitrogen with mortar and pestle. Protein was extracted in buffer (50 mM Tris-HCl buffer, pH 7.5, 150 mM NaCl, 10% v/v glycerol, 0.1% w/v Nonidet P-40) supplemented with 1 mM PMSF, and complete protease inhibitor cocktail (Roche, Indianapolis, IN, USA). The crude extract was filtered through four layers of cheesecloth and centrifuged at 12,000 ×g for 10 min at 4°C. The protein concentration in the supernatant was determined by Bradford assay (Bradford, 1976). Extracts containing similar amounts of total protein were incubated with 500 µl IgG beads (Amersham Biosciences, UK) for 2 h at 4°C with gentle rotation. After centrifugation at 150 g for 3 min at 4°C, the IgG beads were recovered and washed three times with 10 ml of washing buffer (50 mM Tris-HCl buffer pH 7.5, 150 mM NaCl, 10% v/v glycerol, 0.1% w/v Nonidet P-40) and once with 10 ml of cleavage buffer (50 mM Tris-HCl buffer pH 7.5, 150 mM NaCl, 10% v/v glycerol, 0.1% w/v Nonidet P-40, 1 mM DTT). Elution from the IgG beads was performed by 3h incubation with 50 µl (100 units) of 3C protease (Precision protease; Amersham Biosciences, UK) in 2 ml of cleavage buffer at 4°C with gentle rotation. Released protein was recovered in the supernatant

after centrifugation at 150 g for 3 min at 4°C and, after addition of 1 ml cleavage buffer to the beads and re-centrifugation, in the second supernatant; the supernatants were pooled and stored at 4°C. Pooled released protein was loaded onto 0.5 ml of Co-NTA resin (Qiagen, Valencia, CA, USA) packed into a 10-ml disposable polyprep chromatography column (Bio-Rad). To increase the likelihood of binding, the flow-through was loaded again onto the Co-NTA column before washing with 5x 1 ml washing buffer (50 mM Tris-HCl buffer pH 7.5, 150 mM NaCl, 10% v/v glycerol, 0.1% w/v Nonidet P-40). Elution from the Co-NTA was performed using 5x 0.5 ml of imidazole containing buffer (50 mM Tris-HCl buffer pH 7.5, 150 mM NaCl, 10% v/v glycerol, 0.1% w/v Nonidet P-40, and 0.3 M imidazole). Protein in the fractions was bound to 10 µl strataresin (Stratagene company, La Jolla, USA), and separated by SDS-PAGE for subsequent staining with Coomassie Blue and for blotting and detection of the TAP-tagged proteins.

GUS activity detection: Seedlings and different parts of transgenic GUS plants were incubated in a solution containing 1 mM X-Gluc buffer, 0.5 mM potassium ferrocyanide, 0.5 mM potassium ferricyanide, 0.016 mM Triton X-100, 10 mM EDTA, and 100 mM of NaH₂PO₄ and Na₂HPO₄ pH 7.0 at 37°C for 10 h. To visualize the blue precipitate, chlorophyll was removed in 70% v/v ethanol after staining.

Isolation and characterization of homozygous T-DNA insertion lines: Seeds were obtained from the Nottingham *Arabidopsis* Stock Centre (Nottingham, UK). Homozygous *gr-rbp2* mutant plants were isolated from seeds of segregating T3 lines of the T-DNA insertion lines SALK-059714, SALK-13850 (*GR-RBP2*). The insertion in the gene was identified using T-DNA left border primer LBa (5'-TGGTTCACGTAGTGGGCCATCG-3') and specific primers for SALK-059714 (Forward: 5'-TTAGAATGCACCTAATGGGG-3' and Reverse: 5'-CTACTTGTTTTG-GATCAGCCC-3'), SALK-13850 (Forward: 5'-TTA-GAATGCACCTAATGGGG-3' and Reverse: 5'-CCCATCTCGTTGAATTGATG-3').

RESULTS

Phylogenetic analyses of GR-RBP2: To determine the phylogenetic relationship, *GR-RBP2* homologous sequences were retrieved from NCBI and TAIR. Cluster analysis revealed that the protein is identical to the cDNA

product GRP2. It was named earlier in the “plant gene register” section of Plant Physiology by Macknight (1997). Here to avoid confusion we use GR-RBP2 (At4g13850), as named by Lordovic and Bartel (2002). Alignment of the eight GR-RBP paralogs in *Arabidopsis* reveals that the GR-RBPs are conserved in their RRM domains and more divergent in the glycine-rich domain (Fig. 1A). GR-RBP2, GR-RBP3, GR-RBP4, GR-RBP5 and GR-RBP6 have mitochondrial targeting sequences. The closest homologue to GR-RBP2 is GR-RBP4 (Fig. 1B). Alignment of GR-RBP2 with proteins from all organisms revealed that the protein is conserved in both monocotyledonous and dicotyledonous plants. Homologues to GR-RBP2 can be found in other eukaryotes and bacteria (Fig. 1C). Ribosomal protein RPS19 (RPS19) from *Arabidopsis* is close homologue to GR-RBP2. Homology is high in the single RRM sequence but S19 lacks the low complexity Glycine rich region.

Isolation and characterization of knockout mutants of GR-RBP2: Initially, T-DNA insertion lines of GR-RBP2 were considered for functional analysis. Homozygous T-DNA mutant lines were isolated from segregating T3 plants. The knockout phenotype in plants from SALK-059714 is caused by T-DNA insertion in the coding region of *GR-RBP2* (Fig. 2A). Transcript analysis of homozygous lines showed absence of expression of *GR-RBP2* in mutants (Fig. 2B). When wild type and knock out lines are growing on 100 mM trehalose, there was not any difference between them (Fig. 2C). Response of these seedlings to fructose, glucose or sucrose was similar to WT (data not shown). Taken together, *GR-RBP2* mutants were identified but failed to reveal phenotypic of GR-RBP2 with respect to trehalose sensitivity or response to other. Functional redundancy with other members of the multi-gene families of *GR-RBP2* may be one reason for the absence of altered phenotypes.

GR-RBP2 is expressed in all developmental stages: To examine the organ specific expression of *GR-RBP2*, plant lines were transformed that express a transcriptional fusion of GUS using a 2 kb promoter and 5' UTR of *GR-RBP2*. As a control, empty vector of pMDC162 was transformed in WT plants. Strong GUS expression specific to the *GR-RBP2* promoter was detected in the root tips of *Arabidopsis* seedlings (Fig. 3A). Gus activity was observed in adult leaves (Fig. 3B), flowers (Fig. 3C) and siliques (Fig. 3D), Whilst Plants expressing the empty control plasmid did not stain (Fig. 3E, F). The expression pattern of the GUS-fusions was affected by glucose, sucrose or trehalose




Figure 1. *Arabidopsis* GR-RBPs paralogs and ortologs. **A:** Aligned amino acid residues of GR-RBP protein in *Arabidopsis*. The lines indicate conserved motifs in the RRM (RNP1, RNP2) and the glycin-rich domain. **B:** Phylogram derived from alignment of the eight *Arabidopsis* GR-RBP2. The method used is the NJ (Neighbour Joining) method of Saitou and Nei, 1987. **C:** Phylogram derived from GR-RBP2's closest homologues in NCBI. AtGR-RBP2 and AtGR-RBP4, *Arabidopsis thaliana* GR-RBP protein 2 and 4; AtRPS19, *Arabidopsis thaliana* Ribosomal protein S19; PsGR-RBP, *Pisum sativum* GR-RBP protein; MiRNP, *Medicago truncatula* RNA binding protein; StRBP, *Solanum tuberosum* RNA binding protein; BrGR-RBP, *Brassica rapa* GR-RBP protein; Os01g0916660, Os07g0602600 and Os12g0502200, proteins from loci in *Oriza sativa*; PpGR-RBP, *Physcomitrella patens* GR-RBP protein; PgGR-RBP, *Picea glauca* GR-RBP protein; HvGR-RBP, *Hordeum vulgare* GR-RBP.

treatment. These results indicated that the *GR-RBP2* promoter is not responsive to these sugars (data not shown).

Taken together, *GR-RBP2* appeared to be expressed in plants and at all developmental stage; its highest expression was in root tips. *GR-RBP2* expression seems insensitive to trehalose or metabolisable sugars.

GR-RBP2 is likely found in a protein complex in

planta: To identify complexes containing *GR-RBP2* and eventually the factors that interact with *GR-RBP2*, plants were transformed that express a tagged version of *GR-RBP2*. *GR-RBP2* was fused to the Tandem Affinity Tag (TAPa) at the C-terminal end (*GR-RBP2*: TAP); the TAP tag has protein A, myc- and his- epitopes (Rubio *et al.*, 2005). To be able to test if the fusion protein is functional, the CaMV35S promoter was chosen to drive the expression of the tagged pro-




Figure 2. Knock out mutants of *GR-RBP2*. A: Schematic representation of T-DNA insertion in *gr-rbp2* mutant. B: Expression of *GR-RBP2* in knockout lines of *gr-rbp2* (Salk-059714). "WT" wild type seedlings, "mutant" plant lines homozygous for insertions in *GR-RBP2*. *AtACTIN2* expression levels are shown as control. C: Seedlings from WT and lines with *GR-RBP2*-knockouts (*gr-rbp2*) after 14 d on 100 mM of either trehalose or sorbitol under long day conditions at 22°C.

tein: over-expression of a functional protein leads to trehalose resistance. As control plants were transformed with empty vector. In addition, control plant lines were transformed with protein N-terminally tagged and empty plasmid. Plant lines were screened for expression of tagged *GR-RBP2* by western blotting using Anti-myc antibody. This antibody typically detected several protein bands in plants expressing *GR-RBP2:TAP*. But the N-terminally tagged protein or plants with empty vector control did not express any protein (data not shown). The highest molecular weight band corresponds to a protein of 43 kD, which is the expected size for the *GR-RBP2:TAP* fusion before its targeting into the mitochondrion.

Transgenic lines expressing *GR-RBP2:TAP* were also tested for their ability to grow on 100 mM trehalose. Seedling roots of the transgenic lines were not significantly longer than WT. Extract of plants expressing *GR-RBP2:TAP* was separated by gel filtration and the fractions analyzed by western blotting




Figure 3. *GR-RBP2* gene expression and protein-localization. Typical GUS staining in plants containing the translational fusion in A: root tip, B: leaves, C: inflorescence, D: siliques, E and F: in a leaf and siliques of plants transformed with pMDC160 empty vector control. The typical staining pattern shown was found in all three independent transgenic lines tested from the nine lines raised. (Bar = 0.5 cm).

using anti-myc antibody (Fig. 4). Absorption of proteins after separation on the column at 280 nm revealed that the major peak of Rubisco eluted at the size expected from calibration of the system with protein standards (Fig. 4A). Dominant 43 kD precursor protein in crude extracts was detected by Anti-myc antibodies (Fig. 4B, line T). Anti-myc further detected two major processed or degraded forms of *GR-RBP2*: TAP in fractions spanning E6 to E9 that correspond to complexes of approximately 95-100 kD (Fig. 4B, lines E6-E9). A very faint signal was observed in fraction F3 that corresponds to proteins of approximately 45 kD, close to the monomeric size of tagged *GR-RBP2* (Fig. 4C, line F3). Gel-filtration experiments with extracts from control plants did not reveal any components cross-reacting with anti-myc antibody (data not shown). Therefore, *GR-RBP2:TAP* appears to occur in a complex of at least twice its molecular weight (95-100 kD). Tandem affinity purification of the complex was then attempted according to Rubio *et al.* (2005). Preliminary results showed that the procedure allowed enriching a specific set of proteins including protein




Figure 4. GR-RBP2 complex *in planta*. A: Overexpressors of GR-RBP2:TAP on 100 mM trehalose. Seedlings from a line over-expressing *GR-RBP2:TAP* and WT were grown on half strength MS medium with 100 mM trehalose for 2 weeks in long day conditions. B: Sepharose S-300 gel filtration of an extract of plants expressing *GR-RBP2:TAP*. Absorption at 280 nm of proteins after separation on the column: Rubisco, absorbance peak corresponding to rubisco protein; E9, F3 fractions that contain the GR-RBP2 fusion as detected in C. C: Detection of GR-RBP2:TAP by western blotting using anti-myc antibody in fractions E6 to G3 obtained during gel filtration. T, total crude extract; 95 and 45, fractions corresponding to molecular weight 95 and 45 kD, respectively.

corresponding to processed GR-RBP2: TAP (data not shown).

DISCUSSION

In *Arabidopsis* seedlings, exogenously supplied trehalose inhibits root growth and emergence of leaves (Wingler *et al.*, 2000). This growth inhibition is due to T6P accumulation (Schluepmann *et al.*, 2004). Evidence is accumulating that suggests an important regulatory role for T6P. It was shown that T6P deter-

mines carbon utilization in *Arabidopsis* seedlings. However it is not understood how T6P controls carbon utilization and why changes in the steady state level of T6P yields such strong phenotypic changes. In current report we characterized a suppressor of T6P mediating growth arrest using a genetic approach.

GR-RBP2 is a small protein with binding sites for RNA, through its RRM domain, and for protein-protein interaction through its glycine-rich domain. GR-RBP2 is a protein of likely prokaryotic origin that is part of a protein complex. *GR-RBP2* expression is found throughout the plant and is not responsive to

metabolisable sugars such as glucose, fructose or sucrose. *GR-RBP2* expression also does not respond to 100 mM trehalose. A T-DNA insertion mutant of *GR-RBP2* was identified in the T-DNA insertion line SALK-059714. These mutants did not show different phenotype from WT, when grown on different carbon sources. The data obtained did not reveal a function for *GR-RBP2* with respect to growth in relation to the carbon metabolism.

Gus expression data revealed that *GR-RBP2* is highly expressed in young seedlings, flowers and root tips. High expression of *GR-RBP2* in young seedlings suggests that may have a capability to play functions in RNA metabolism in actively proliferating organs (Kwak *et al.*, 2005). Transcript analysis data were in accordance with the data from gene expression databases such as Genevestigator (Zimmermann *et al.*, 2004), suggesting that the 2kb promoter sequence contains the regulatory sequences driving *GR-RBP2* expression *in planta*.

TAPa-tagged fusion and gel-filtration data showed that *GR-RBP2* is likely in a complex of 95-100 kD in plants. A major lower molecular weight bands may correspond to the 40 kD processed protein that had been imported in the mitochondrion or to degraded products. Further work is however needed to analyze proteins co-purifying with *GR-RBP2*:TAP.

GR-RBPs are involved in abiotic stress response and their functions were recently indicated (Kim *et al.*, 2008; Kim *et al.*, 2007; Kwak *et al.*, 2005). Previous studies showed that *GR-RBP* 1, 2, 3, 4, 7 and 8 expressions are increased by cold stress (kim *et al.*, 2007; Kwak *et al.*, 2005). Meanwhile, *GR-RBP1* expression was strongly induced by dehydration and salt stress (Kwak *et al.*, 2005).

An improvement of cold tolerance was noted in over-expressing *GR-RBP2* plants (Kim *et al.*, 2007; Kwak *et al.*, 2005; Vernel *et al.*, 2002). It was also demonstrated that *Arabidopsis GR-RBP2* is able to suppress the cold sensitivity of *E. coli* at low temperatures, which suggests that *GR-RBP2* exhibits an RNA chaperone activity during cold adaptation. Plant over-expressing *GR-RBP2* showed induced catalase and peroxidase but decreased superoxide dismutase activities. The closest homologue to *GR-RBP2*, *GR-RBP4*, is unable to complement *E. coli* BX04. Accordingly, the RNA chaperone activity of *GR-RBP2* is specific (Kim *et al.*, 2007). This specificity may arise from complex formation with other factors (Fig. 4). Further characterization of the proteins in the complex with *GR-RBP2* will be necessary to understand *GR-RBP2*

function. Possibly, *GR-RBP2* over-expression leads to the growth on trehalose due to its stress protection function. T6P accumulation might generate metabolic stress that can be alleviated by *GR-RBP2*. In *GR-RBP2* over-expressors starch accumulation on trehalose is similar to WT. Therefore, *GR-RBP2* is an indirect target of T6P. *GR-RBP2* likely affects process downstream of T6P.

Kwak *et al.* (2005) suggested that *GR-RBPs* originate from cyanobacteria where they replace the function of cold-shock proteins as found in other bacteria such as in *E. coli*. *GR-RBPs* are also present in mammals, e.g. the cold induced RNA binding proteins (Nishiyama *et al.*, 1997). Cold induced RNA binding proteins (CIRP) of the mouse is involved in growth inhibition under reduced temperature (Nishiyama *et al.*, 1997). Over-expression of CIRP leads to prolongation of the G1 phase of the cell cycle. Immuno-fluorescence studies located CIRP in the nucleus, unlike *GR-RBP2* that is located in mitochondria (Fig. 3). Surprisingly the products of this gene are found in mitochondria whilst T6P biosynthesis is thought to occur in the cytosol (TAIR annotation of TPS genes, March, 2007). None of the proteins of T6P metabolism in *Arabidopsis* have a mitochondrial targeting sequence (Schluepmann *et al.*, 2004; Leyman *et al.*, 2001) and none have been found associated with mitochondrial proteome (Kruft *et al.*, 2001). The link with reactions in mitochondria through *GR-RBP2* is novel. The function of *GR-RBP2* as an RNA chaperone (Kim *et al.*, 2007) suggests that it may alter transcription of mitochondrial proteins that affect T6P targets in the cytosol or chloroplast.

In conclusion, the present work provides new information that increases our knowledge about the roles of *GR-RBP2* in response of plants to trehalose. *GR-RBP2* likely exerts its effect upon growth by changing mitochondrial metabolism. Taken together, *GR-RBP2* links mitochondrial stress adaptation and T6P metabolism and this link may begin to explain how trehalose metabolism mediates stress protection in plants. This protein is likely modulating a component down stream of T6P.

Acknowledgments

We are greatly thankful to Dr. Viola Willemsen for kindly providing pDONR and N-TAPa TAG vectors, Mark de Jong for providing pMDC162 and pMDC107 vectors, the Nottingham *Arabidopsis* Stock Center (NASC) for providing seeds and Frits Kindt and Ronald Leito for help with confocal microscope and photographs.

References

- Aghdasi M, Schluepmann H (2009). Cloning and expression analysis of two photosynthetic genes, *PSI-H* and *LHCBI*, under trehalose feeding conditions in *Arabidopsis* seedlings. *Iran J Biotechnol.* 7: 179-187.
- Aghdasi M (2007). Analysis of T6P control over carbon allocation and growth in plants. Utrecht University, Utrecht. PP: 51-67 (*Ph.D thesis*).
- Bradford MM (1976). A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 72: 248-254.
- Carpenter CD, Kreps JA, Simon AE (1994). Genes encoding glycine-rich *Arabidopsis thaliana* proteins with RNA-binding motifs are influenced by cold treatment and endogenous circadian rhythm. *Plant Physiol.* 104: 1015-1025.
- Clough SJ, Bent AF (1998). Floral dipping: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16:735-743.
- Condit CM, Meagher RB (1987). Expression of a gene encoding a glycine rich protein in *Petunia*. *Mol Cell Biol.* 7: 4273-4279.
- Curtis MD, Grossniklaus U (2003). A gateway cloning vector set for high-throughput functional analysis of genes in plants. *Plant Physiol.* 133: 462-469.
- De Oliverira DE, Seurinck J, Inze D, van montagu M, Botterman J (1990). Differential expression of five *Arabidopsis* genes encoding glycine-rich proteins. *Plant Cell.* 2: 427-436.
- Elbein AD, Pan YT, Pastuzak I, Carroll D (2003). New insights on trehalose: a multifunctional molecule. *Glycobiology* 13: 17R-27R.
- Eastmond PJ, Van Dijken AJ, Speiman M, Kerr A, Tissier AF, Dichinson HG, Jones JD, Smeekens SC, Graham IA (2002). Trehalose-6-phosphate synthase 1 which catalyses the first step in trehalose synthesis is essential for *Arabidopsis* embryo maturation. *Plant J.* 29: 223-235.
- Fang RX, Pang Z, Gao DM, Mang KQ, Chua NH (1991). cDNA sequence of a virus-inducible, Glycine-rich proteins gene from rice. *Plant Mol Biol.* 17: 1255-1257.
- Fritzius T, Aechbacher R, Wiemken A, Wingler A (2001). Identification of ApL3 expression by trehalose complements the starch-deficient *Arabidopsis* mutant *adg2-1* lacking ApL1, the large subunit of ADP-glucose pyrophosphorylase. *Plant Physiol.* 126: 883-889.
- Fusaro AF, Bocca SN, Ramos RL, Barroco RM, Magioli C, Jorge VC, Coutinho TC, Rangel-Lima CM, De Rycke R, Inze D, Engler G, Sachetto-Martins G (2007). AtGRP2, a cold-induced nucleo-cytoplasmic RNA-binding protein, has a role in flower and seed development. *Plant J.* 225: 1339-1351.
- Goddijn O, Smeekens S (1998). Sensing trehalose biosynthesis in plants. *Plant J.* 14: 143-146.
- Kim JY, Park SJ, Jung C, Ann S, Goh CH, Cho K, Hano O, Kang H (2007). Functional characterization of a glycine-rich RNA binding protein2 in *Arabidopsis thaliana* under abiotic stress conditions. *Plant J.* 50: 439-451.
- Kim JS, Hung HJ, Lee HJ, Kim KA, Goh CH, Woo Y, Oh SH, Han YS, Kang H (2008). Glycine-rich RNA binding protein7 affects abiotic and closing in *Arabidopsis thaliana*. *Plant J.* 55: 455-466.
- Kolbe A, Tiessen A, Schluepmann H, Paul M, Ulrich S, Geigenberger P (2005). Trehalose 6-phosphate regulates starch synthesis via posttranslational redox activation of ADP-glucose pyrophosphorylase. *Proc Natl Acad Sci U.S.A.* 102: 11118-11123.
- Kruft V, Eubel H, Jansch L, Werhahn W, Braun HP (2001). Proteomic approach to identify novel mitochondrial proteins in *Arabidopsis*. *Plant Physiol.* 127: 1694-1710.
- Kwak KJ, Kim YO, Kang H (2005). Characterization of transgenic *Arabidopsis* plants overexpressing GR-RBP2 under high salinity, dehydration or cold stress. *J Exp Bot.* 421: 3007-3016.
- Leclere J, Bartel B (2001). A library of *Arabidopsis* 35S-cDNA lines for identifying novel mutants. *Plant Mol Biol.* 46: 695-703.
- Leyman B, Van Dijk P, Thevelein JM (2001). An unexpected plethora of trehalose biosynthesis genes in *Arabidopsis thaliana*. *Trends Plant Sci.* 6: 510-513.
- Lorkovic ZJ, Barta A (2002). Genome analysis: RNA recognition motif (RRM) and K homology (KH) domain RNA-binding proteins from flower plant *Arabidopsis thaliana*. *Nucleic Acids Res.* 30: 623-635.
- Lorkovic ZJ (2009). Role of plant RNA-binding proteins in development, stress response and genome organization. *Cell* 14: 229-237.
- Macknight R (1997). FCA, a gene controlling flowering time in *Arabidopsis*, encodes a protein containing RNA binding domain. *Cell* 89: 737-745.
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant.* 15: 473-479.
- Nishiyama H, Higashitsuji H, Yokoi H, Itoh K, Danno S, Matsuda T, Fujita J (1997). Cloning and characterization of human CIRP (cold inducible RNA-binding protein) cDNA and chromosomal assignment of the gene. *Gene.* 204: 115-120.
- Nomata T, Kabeya Y, Sato M (2004). Cloning and characterization of glycine-rich RNA-binding proteins cDNAs in the moss *Physcomitrella patens*. *Plant Cell Physiol.* 45: 48-56.
- Pramanic MH, Imai R (2005). Functional identification of a trehalose-6-phosphate phosphatase gene that is involved in transient induction of trehalose biosynthesis during chilling stress in rice. *Plant Mol Biol.* 58: 751-762.
- Paul M, Lucia F, Primavesi F, Deveraj J, Zhang Y (2008). Trehalose metabolism and signaling. *Ann Rev Plant Biol.* 59: 417-441.
- Pellny TK, Ghannoum O, Conroy JP, Schluepmann H, Smeekens S, Andralojc J, Krause KP, Goddijn O, Paul MJ (2004). Genetic Modification of photosynthesis with *E. coli* genes for trehalose synthesis. *Plant Biotechnol J.* 2: 71-82.
- Ramon M, Rolland F, Thevelein JM, Van Dijk P, Leyman B (2007). ABI4 mediates the effects of exogenous trehalose on *Arabidopsis* growth and starch breakdown. *Plant Mol Biol.* 63: 195-206.
- Rubio V, Shen Y, Saijo Y, Liu Y, Gusmaroli G, Dinesh-Kumar SP,

- Deng XW (2005). An alternative tandem affinity purification strategy applied to Arabidopsis protein complex isolation. *Plant J.* 41: 767-778.
- Rolland F, Baena-Gonzales E, Sheen J (2006). Sugar sensing and signaling in plants: conserved and novel mechanisms. *Annu Rev Plant Biol.* 57: 675-709.
- Salinas-Jose J, Sanchez-Serrato J (2006). Method in biology: Arabidopsis protocols. *Humana Press*. PP: 216-217.
- Schlupepmann H, Pellny T, Van Dijken A, Smeekens S, Paul MJ (2003). Trehalose-6-phosphate is indispensable for carbohydrate utilization and growth in Arabidopsis Thaliana. *Proc Natl Acad Sci USA.* 100: 6849-6854.
- Schlupepmann H, Van Dijken A, Aghdasi M, Wobbes B, Paul M, Smeekens S (2004). Trehalose mediated growth inhibition of Arabidopsis seedlings is due to trehalose-6-phosphate accumulation. *Plant physiol.* 135: 879-890.
- Vandesteene L, Ramon M, Le Roy K, Van Dijk P, Rolland F (2010). A single active Trehalose-6-p synthase (TPS) and a family of putative regulatory TPS-like proteins in Arabidopsis. *Molecular Plant advance.* 2: 1-14.
- Vermel M, guermann B, Delage L, Grienberger JM, Marechal-Drouar L, Gualberto JM (2002). A family of RRM-type RNA-binding proteins specific to plant mitochondria. *Proc Natl Acad Sci USA.* 99: 5866-5871.
- Wingler A, Fritzius T, Wiemken A, Boller T, Aeschbacher RA (2000). Trehalose induces the ADP-glucose pyrophosphorylase gene, ApL3, and starch synthesis in Arabidopsis. *Plant Physiol.* 124: 105-114.
- Zimmermann P, Hirsch-hoffmann M, Hennig L, Grissem W (2004). Geninvestigator Arabidopsis microarray database and analysis toolbox. *Plant physiol.* 136: 2621-2632.