



In vitro versus *In vivo*: Development-, Apoptosis-, and Implantation-Related Gene Expression in Mouse Blastocyst

Forough Mahdavinezhad¹, Parinaz Kazemi^{1,2}, Parisa Fathalizadeh¹, Fatemeh Sarmadi¹, Ehsan Hashemi¹, Hadi Hajarian³, Mojtaba Dashtizad^{1,*}

¹ Embryo Biotechnology Laboratory (Embio Lab), Department of Animal Biotechnology, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

² McGill University Research Institute of the McGill University Health Centre, Montreal, Quebec, Canada

³ Department of Animal Science, Razi University, Kermanshah, Iran

* Corresponding author: Mojtaba Dashtizad, Department of Animal Biotechnology, National Institute of Genetic Engineering and Biotechnology, Tehran, Iran. Tel: +98-21-44787323; Fax: +98-21-44780395; E-mail: dashtizad@nigeb.ac.ir

Background: While mammalian embryos can adapt to their environments, their sensitivity overshadows their adaptability in suboptimal *in vitro* conditions. Therefore, the environment in which the gametes are fertilized or to which the embryo is exposed can greatly affect the quality of the embryo and consequently its implantation potential.

Objectives: Since providing an optimal culture condition needs a deep understanding of the environmental effects, and regarding the fact that normal morphology fails to be a reliable indicator of natural embryo development, the current study aimed at comparing *in vivo*- and *in vitro*-derived blastocysts at the molecular level.

Materials and Methods: *In vivo* and *in vitro* mouse blastocysts were obtained by flushing the uterine horns and *in vitro* fertilization/culture, respectively. Normal blastocysts of both groups were evaluated in terms of hatching rate and expression of three lineage-differentiation-, apoptosis-, and implantation-related genes.

Results: The hatching rate was lower in *In vitro* fertilization (IVF)-produced blastocysts in comparison with that of the *in vivo* counterparts. More importantly, the study results indicated significant changes in the expression levels of eight out of ten selected genes, especially *Mmp-9* (about -10.7-fold). The expression of *Mmp-9* in trophoblast cells is required for successful implantation and trophoblast invasion.

Conclusions: The current study, in addition to confirming that the altered gene expression pattern of *in vitro*-produced embryos resulted in normal morphology, provided a possible reason for lower implantation rate of *in vitro*-produced blastocysts regarding the *Mmp-9* expression.

Keywords: Gene Expression; Fertilization *in vitro*; Matrix Metalloproteinase 9

1. Background

In vitro culture (IVC) of preimplantation embryo is undoubtedly an essential step in assisted reproductive technologies such as *in vitro* fertilization (IVF) and intracytoplasmic sperm injection, as well as embryonic stem cell studies (1, 2). Mammalian preimplantation embryo development period extends from fertilization of mature oocyte to implantation of the late blastocyst (3). During this interval, many important events including maternal to zygotic genome transition, compaction, first three lineage differentiation, and blastocyst formation occur. These events seem to be affected by gamete quality and culture conditions (4) such as culture media components, pH, osmolarity, temperature, oxygen tension, and humidity of incubator (5-7).

Despite the significant improvements in the quality of culture media (8, 9), it seems that none of the available ones can fully mimic the physiological conditions of the female tract (10). In this regard, many studies in different species show that environmental stresses can cause serious changes in morphology and cell metabolism (11-17), affecting the quality of the embryo and consequently its implantation potential (18, 19). *In vitro*-produced embryos have comparably higher lipid accumulation in the cytoplasm (20), more fragile zona pellucida (21), higher chromosomal abnormality (22), smaller nucleoplasmic rate (23), and reduced total number of cells (24). The origin of these cellular and sub-cellular differences between *in vivo*- and *in vitro*-derived embryos can be traced to gene expression alterations (25). In fact, although embryos can adapt to

an artificial environment, suboptimal conditions might immediately affect gene expression. However, the consequences might not be evident until later stages of embryo development, fetus phase or even postnatal period (26, 27). The first sign of these invisible effects is low rate of implantation even after transferring a morphologically normal embryo (18, 28, 29). Since the normal morphology of an embryo fails to guarantee its high quality, *in vitro* culture conditions are favored to be further assessed at molecular levels. In fact, identifying genes with different expression patterns in *in vitro* environment may facilitate providing an optimal culture condition with more appropriate factors.

2. Objectives

The current study aimed at comparing the hatching rate and the expression of selected genes related to developmental potential, implantation ability, and apoptosis between *in vivo*- and *in vitro*-derived mouse blastocysts. More specifically, the study examined relative levels of caudal-type homeobox 2 (*Cdx2*) and eomesodermin (*Eomes*), both involved in trophoblast differentiation (30); pluripotency-sustaining factors POU class 5 homeobox 1 transcription factor (*Pou5f1*, formerly *Oct4*) and *Nanog* homeobox protein (*Nanog*), essential for inner cell mass formation (31); GATA-binding factor 6 (*Gata6*), the growth receptor bound protein 2 (*Grb2*)-RAS-mitogen-activated protein (MAP) kinase signaling (32), drives differentiation towards the primitive endoderm; and matrix metalloproteinase 9 (*Mmp-9*), associated with extracellular matrix degradation during implantation process (33). Furthermore, transformation-related protein 53 (*Trp53*), which plays a critical role in the initiation of apoptosis and its downstream target genes, *Bcl2*-associated X (*Bax*), and B-cell lymphoma 2 (*Bcl2*), were chosen as proxies to understand how *in vitro* culture stresses the embryo (34, 35).

3. Materials and Methods

All mice were housed in the National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran. Facility and procedures of using these mice were reviewed and approved by the NIGEB Institutional Animal Care and Use Committee. Studies were performed in accordance with the guidelines for the Care and Use of Laboratory Animals (IR.NIGEB.EC.1394.8.10.A).

3.1. *In vivo* and *in vitro* Blastocyst Production

Female NMRI mice (6–8 weeks old) were induced to superovulate by intraperitoneal injection of 7 IU equine chorionic gonadotropin (eCG; Folligon, Intervet, Spain) followed by 7 IU human chorionic gonadotropin 48 hours later (hCG; Pregnyl, Daroupakhsh, Tehran, Iran). For *in vivo* blastocyst collection, 21 superovulated female mice (seven in each replicate) were placed overnight with fertile males from the same strain (1:1).

The success of mating was confirmed the next morning by checking the vaginal plug. The date of plug detection considered E0.5. Mated females were sacrificed by cervical dislocation and blastocysts were obtained at E4.5 by flushing each uterine horn with HEPES buffered M2 medium (M7167; Sigma-Aldrich, St. Louis, MO, USA) and then cultured in KSOM (MR-121-L; Millipore, Billerica, MA, USA) medium at 37°C in a highly humidified atmosphere containing 5% CO₂ under a mineral oil overlay (M5310; Sigma-Aldrich).

For *in vitro* blastocyst production, 12–14 hours after hCG injection, 42 superovulated female mice (13–15 for each replicate) were sacrificed by cervical dislocation. Cumulus-oocyte complexes (COCs) were obtained from the ampulla of the oviducts and washed into the M2 medium. After washing, groups of 10–15 COCs were placed in 50 µL droplet of human tubal fluid (HTF) medium (36) supplemented with 6 mg.mL⁻¹ BSA (A6003; Sigma-Aldrich), 36.3 mg.mL⁻¹ sodium pyruvate (P5280; Sigma-Aldrich) and 30.7 mg.mL⁻¹ glutathione (G4251; Sigma-Aldrich). The sperm was obtained by mincing the vasa deferentia and each cauda epididymides male NMRI mice (8–10 weeks old) into HTF medium. The sperm dish was placed in the incubator under conditions described earlier for 30 minutes to allow the sperm swim out. Capacitated sperms were then added to HTF droplets containing oocytes for a final concentration of 1×10⁶/mL. After four hours, presumptive zygotes were washed and cultured in KSOM medium to the blastocyst stage. Finally, morphologically normal expanded blastocysts with thinning zona were selected for the next experiments.

3.2. RNA Isolation and Real-time RT-PCR

Total RNA was isolated from 100 blastocysts for three replicates in each group (32–35 blastocysts in each replicate), using RNase plus Micro Kit (Qiagen, Valencia, CA, USA) in accordance with the manufacturer's protocol. The extracted RNA was converted into cDNA by AccuPower[®] RocketScript[™] RT PreMix kit (Bioneer, Daejeon, Korea) using random hexamer primers. The PCR reaction was prepared at the final volume of 15 µL by mixing 7.5 µL of 2X SYBR Green PCR Master Mix (25344; Intron, Seongnam-Si, Gyeonggi-do, Korea), 3 µL of cDNA, and 0.2 µM of each primer. Subsequently, real-time PCR was performed with a ABI System (Applied Biosystems StepOne, Foster City, CA, USA) under the following thermal conditions: 95°C for two minutes, 40 cycles of 95°C for 10 seconds, and 58°C for 30 seconds. After 40 cycles, melting curves were analyzed to confirm the specificity of PCR products. Finally, relative expression of each gene was determined by the 2^{-ΔΔCt} method, using *B2m* (beta-2-microglobulin) as the reference gene (37, 38). The sequences of the used primers (synthesized by TaqCopenhagen, Copenhagen, Denmark) are listed in **Table 1**.

Table 1: Primer sequences for RT-PCR analysis

Gene	Accession Number	Primer Sequence (5' to 3')	Product Size
<i>Trp53</i>	NM_011640	F: TGGAGGAGTCACAGTCGGAT R: CGTCCATGCAGTGAGGTGAT	117
<i>Bax</i>	NM_007527	F: GGTGCTCAAGGCCCTGTG R: GGTCCCGAAGTAGGAGAGGAG	142
<i>Bcl2</i>	NM_009741	F: CTTCGCAGAGATGTCCAGTCAG R: CCACAATCCTCCCCAGTTCAC	113
<i>Pou5f1</i>	NM_013633	F: AGCATTGAGAACCCTGTGAGG R: TCGAACACATCCTTCTCTAGC	120
<i>Nanog</i>	NM_028016	F: GCCTCCAGCAGATGCAAGAA R: GGTGCTGAGCCCTTCTGAAT	154
<i>Cdx2</i>	NM_007673	F: GGAGGAAAAGTGAGCTGGCTG R: CTCTCCTTGGCTCTGCGGTT	85
<i>Eomes</i>	NM_010136	F: CCCACTGGATGAGGCAGGAGA R: CCACACCGTCTCTGTCACTT	153
<i>Gata6</i>	NM_010258	F: CAGGGGTAGGGGCATCAGTG R: GCAGGGGAGGACAGACTGAC	118
<i>Grb2</i>	NM_008163	F: CACGGGTGGCATGTGTGTC R: AAGCAGGGGGAAGGGAATC	101
<i>Mmp-9</i>	NM_013599	F: GCAAAGGCGTCGTGATCC R: TGCCGTCCTTATCGTAGTCAG	162
<i>B2m</i>	NM_009735	F: CCTGGTCTTCTGGTGCTTGT R: GCAGTTCAGTATGTTCCGGCTTC	118

F: forward; R: reverse.

3.3. Experimental Design

In order to investigate the effects of *in vitro* condition on embryo quality, after evaluation of blastocyst formation rate, the number of hatched blastocysts in IVF treatment and *in vivo* control groups were counted and compared; the hatching rate was measured as follow: number of hatching, hatched blastocysts/ total blastocyst. After that, the blastocysts were separately stored at -80 °C until RNA extraction. Finally, the transcript levels of the desired genes were quantified in both groups by real-time PCR.

3.4. Statistical Analysis

All experiments were repeated at least three times and results were expressed as mean ± SD. The Student t-test

was employed to analyze the experimental data. The SPSS version 16.0 was used for this analysis and P-value of <0.05 was considered statistically significant.

4. Results

4.1. Effect of In vitro Fertilization and Culture on Blastocyst Formation

To assess the impact of *in vitro* fertilization and culture media on developmental competence of embryo during preimplantation stages, the oocytes were fertilized and cultured *in vitro* for 96 hours. As shown in **Table 2**, a total of 351 two-cell embryos, 277 morulae, and 264 blastocysts (64.2%, 50.55%, and 48.17%, respectively) were produced from *in vitro* fertilization of 548 oocytes.

Table 2. Development of *in vitro* fertilized mouse oocytes

Group	Oocytes, N	2-Cell Embryo, N (%)	Morula, N (%)	Blastocyst, N (%)
<i>In vitro</i> fertilization	548	351 (64.2 ± 2.79)	277 (50.55 ± 2.56)	264 (48.17 ± 1.77)

4.2. Effect of In vitro Fertilization and Embryo Culture on Hatching Rate of Embryos

To further investigate the effects of *in vitro* condition, morphologically normal expanded blastocysts of IVF treatment and *in vivo* control groups were cultured for further 12 hours and analyzed in terms of hatching rate. As shown in **Figure 1**, the percentage of hatched IVF-produced embryos (62.8%) was significantly ($P < 0.05$) lower than that of their *in vivo* (68.47%) counterparts.

4.3. Effect of IVF and Embryo Culture on Developmentally Important Genes

The relative expression of all genes is presented in **Figure 2**. The results revealed that the expression levels of pluripotency genes were significantly ($P < 0.05$)

higher in IVF-produced blastocysts than the fresh ones (4.49- and 1.47-fold, for *Nanog* and *Pou5f1* respectively). However the expression of *Cdx2*, *Gata6*, *Grb2*, and *Mmp9* showed a significant decrease (-5.58-, -2.71-, -5.46-, and -10.77-fold, respectively) in the *in vitro* group. Although the expressions of *Bax* and *Bcl2* were comparatively lower (-2.48-, and -3.32-fold, respectively) in the *in vitro* embryos, the differences in the *Bax:Bcl2* ratio (1.34-fold; $P=0.213$) was not statistically significant. The expression of *Eomes* (1.26-fold, $P=0.088$) and *Trp53* (1.03-fold; $P=0.306$) was similar in both experimental groups.




Figure 1. Hatching rates (%) of *in vivo* and *in vitro* produced blastocysts. ^{a,b}, different letters in the bars indicate statistically significant differences between the experimental groups ($P < 0.05$).




Figure 2. Relative expression of selected genes related to (A) pluripotency, (B) the trophoctoderm lineage, (C) the primitive endoderm, (D) apoptosis, and (E) implantation. Statistically significant differences are indicated by ^{a,b} ($P < 0.05$).

5. Discussion

The current study aimed at assessing the impacts of IVF and IVC on the quality of embryos. According to obtained results, hatching rate was significantly lower in the IVF-derived embryos than in the *in vivo* control. Although a correct hatching process occurs normally in a good quality blastocyst, it is shown that *in vitro* conditions cause zona hardening, and disturb the hatching ability of a produced embryo(39). Furthermore, the success of *in vitro* hatching is dependent on a sufficiently high number of embryonic cells(40) as well as trophoctodermal lysins (41). Schiewe *et al.*, reported that culture conditions may

reduce the production of intrinsic embryonic lysin that promotes hatching(42).

To further elucidate how *in vitro* condition might affect the embryo quality, relative expression levels of *Mmp-9*, which is also referred to as gelatinase B, was examined (43). The results of the current study (Fig. 2) indicated a significantly lower expression level of *Mmp-9* in *in vitro*-produced mouse blastocysts compared with the *in vivo*-produced ones. The expression of *Mmp-9* by trophoblast cells, which possesses basement membrane-degrading proteolytic activity, is required for successful implantation and trophoblast invasion (43, 44). It seems

that *in vitro* culture condition, through a reduction of hatching rate and *Mmp-9* expression, can be related to low implantation rate reported in previous studies (45, 46).

Shortly before implantation, the late blastocyst contains three distinct cell types: epiblast, which forms the future embryo, trophoctoderm, which gives rise to the prospective placenta, and primitive endoderm, which forms the yolk sac (30, 47). In the current study, IVF and IVC significantly increased the expression levels of *Pou5f1* and *Nanog* in the mouse blastocyst. The obtained results were in agreement with those of Henderson *et al.* (48), observing enhanced expression of *Pou5f1* and *Nanog* in the *in vitro*-produced rabbit blastocyst. Purpera *et al.*, also reported a significant upregulation of *Pou5f1* mRNA in *in vitro* bovine blastocysts compared with their *in vivo* counterparts (49). *Pou5f1* and *Nanog* are the most critical transcription factors for creating and maintaining the pluripotency of inner cell mass, epiblast, and embryonic stem cells, as well as the first two lineage differentiation in blastocyst (50). According to the role of these key factors, changing their expression levels can presumably interfere with normal differentiation and developmental competence of embryo (15). In the current study, the expression of *Cdx2* was significantly lower in IVF-produced embryos, while no difference was found about *Eomes*. In contrast, Giritharan *et al.*, (24) reported a decrease (about -2-fold) in *Eomes* expression in *in vitro*-produced mouse blastocyst. *Cdx2* and *Eomes* are two important factors required for trophoctoderm differentiation and development (51). Furthermore, *Cdx2* and *Pou5f1* have mutual inhibitory activity, where *Cdx2* inhibits *Pou5f1* in trophoctoderm cells, and *Pou5f1* inhibits expression of *Cdx2* in the inner cell mass (52, 53). In the current study, reduction in expression of *Cdx2* and an elevation in the expression of the *Pou5f1* show this antagonistic regulation, which in this case can direct cell differentiating toward the inner cell mass (54). Low expression of *Cdx2* was in accordance with that of the study by Giritharan *et al.* (24), in which IVF reduced the trophoctoderm cell numbers. Both of these reductions (in cell number and gene expression) can influence trophoctoderm-specific downstream developmental events such as trophoblast development, implantation, mesoderm, and placenta formation, and therefore, would perturb normal embryo development (51, 55). The current study results revealed that IVF and IVC cause a reduction in the expression of *Gata6* and *Grb2*. *Grb2* and *Gata6* are two essential transcription factors to differentiate and form primitive endoderm in blastocyst (56). According to the fact that the differentiation of primitive endoderm and epiblast occurs due to antagonistic effect of *Gata6* and *Nanog* factors, a reduction in *Gata6* and increase in *Nanog* expression may reinforce leading to the epiblast population, which therefore can perturb the yolk sac

formation and post-implantation embryo development (30, 54).

Programmed cell death or apoptosis, initiating cell death under suboptimal conditions, plays a vital role in the development of the embryo (57). The current study results showed that although the expression levels of *Bax* and *Bcl2* decreased in the *in vitro* group, no significant differences were observed in either the expression of *Trp53* or in the *Bax:Bcl2* ratio between *in vitro* blastocysts and their *in vivo* counterparts. The *Trp53* is a transcription factor that in response to stress signals activates many downstream target genes including *Bax*, which can overcome the anti-apoptotic effects of *Bcl2* and accelerated cell death (35). Thus, the fate of a cell in response to stress can be influenced by *Trp53*, which regulates the ratio of *Bax: Bcl2* protein level (58). Since the ratio of *Bax:Bcl2* is a reliable parameter to predict the tendency of embryo towards survival or apoptosis (34), the current study results indicated a normal *in vitro* condition in terms of apoptosis.

6. Conclusions

The collected data gave further evidence supporting the effects of *in vitro* culture on expression levels of developmental and apoptosis important genes. The expressions of pluripotency genes were significantly higher in IVF-produced blastocysts; whereas, *Cdx2*, *Gata6*, and *Grb2* showed a significant decrease in the *in vitro* group compared with the fresh ones. In addition, the current study provided a possible explanation for the lower implantation rate of *in vitro*-produced blastocysts compared with those of the *in vivo* origin, regarding a sharp decrease in *Mmp-9* expression level. However, further investigations are required to clarify whether returning to normal expression levels of *Mmp-9* through providing an appropriate environment would improve the implantation rate.

Acknowledgments

Authors wish to thank the National Institute of Genetic Engineering and Biotechnology (NIGEB) for providing the facilities to conduct the study.

Financial Support

This work was financially supported by the Iran National Science Foundation (INSF); grant number: 93022945.

Conflict of interest

The authors declared no conflict of interest.

References

1. Calle A, Fernandez-Gonzalez R, Ramos-Ibeas P, Laguna-Barraza R, Perez-Cerezales S, Bermejo-Alvarez P, *et al.* Long-term and transgenerational effects of *in vitro* culture on mouse embryos. *Theriogenology*.

- 2012;77(4):785-793. doi: 10.1016/j.theriogenology.2011.07.016 pmid: 21855990
2. Calle A, Miranda A, Fernandez-Gonzalez R, Pericuesta E, Laguna R, Gutierrez-Adan A. Male mice produced by *in vitro* culture have reduced fertility and transmit organomegaly and glucose intolerance to their male offspring. *Biol Reprod.* 2012;87(2):34. doi: 10.1095/biolreprod.112.100743 pmid: 22649070
 3. Gasperowicz M, Natale DR. Establishing three blastocyst lineages--then what? *Biol Reprod.* 2011;84(4):621-630. doi: 10.1095/biolreprod.110.085209 pmid: 21123814
 4. Lonergan P, Rizos D, Kanka J, Nemcova L, Mbaye AM, Kingston M, et al. Temporal sensitivity of bovine embryos to culture environment after fertilization and the implications for blastocyst quality. *Reproduction.* 2003;126(3):337-346. pmid: 12968941
 5. Johnston LA, Donoghue AM, O'Brien SJ, Wildt DE. Influence of temperature and gas atmosphere on *in vitro* fertilization and embryo development in domestic cats. *J Reprod Fertil.* 1991;92(2):377-382. pmid: 1909365
 6. Rinaudo PF, Giritharan G, Talbi S, Dobson AT, Schultz RM. Effects of oxygen tension on gene expression in preimplantation mouse embryos. *Fertil Steril.* 2006;86(4 Suppl):1252-1265, 1265 e1251-1236. doi: 10.1016/j.fertnstert.2006.05.017 pmid: 17008149
 7. Camargo LSdA, Viana JHM, Sá WFd, Ferreira AdM, Ramos AdA, Vale Filho V. Factors influencing *in vitro* embryo production. *Anim Reprod.* 2006;3(1):19-28.
 8. Summers MC, McGinnis LK, Lawitts JA, Biggers JD. Mouse embryo development following IVF in media containing either L-glutamine or glycyl-L-glutamine. *Hum Reprod.* 2005;20(5):1364-1371. doi: 10.1093/humrep/deh756 pmid: 15705624
 9. Quea G, Romero K, Garcia-Velasco JA. Extended embryo culture to increase implantation rate. *Reprod Biomed Online.* 2007;14(3):375-383. doi: 10.1016/s1472-6483(10)60882-6 pmid: 17359595
 10. Aviles M, Gutierrez-Adan A, Coy P. Oviductal secretions: will they be key factors for the future ARTs? *Mol Hum Reprod.* 2010;16(12):896-906. doi: 10.1093/molehr/gaq056 pmid: 20584881
 11. Doherty AS, Mann MR, Tremblay KD, Bartolomei MS, Schultz RM. Differential effects of culture on imprinted H19 expression in the preimplantation mouse embryo. *Biol Reprod.* 2000;62(6):1526-1535. doi: 10.1095/biolreprod62.6.1526 pmid: 10819752
 12. de Oliveira AT, Lopes RF, Rodrigues JL. Gene expression and developmental competence of bovine embryos produced *in vitro* under varying embryo density conditions. *Theriogenology.* 2005;64(7):1559-1572. doi: 10.1016/j.theriogenology.2005.03.019 pmid: 15878193
 13. Niemann H, Wrenzycki C. Alterations of expression of developmentally important genes in preimplantation bovine embryos by *in vitro* culture conditions: implications for subsequent development. *Theriogenology.* 2000;53(1):21-34. doi: 10.1016/S0093-691X(99)00237-X pmid: 10735059
 14. Sananmuang T, Tharasanit T, Nguyen C, Phutikanit N, Techakumphu M. Culture medium and embryo density influence on developmental competence and gene expression of cat embryos. *Theriogenology.* 2011;75(9):1708-1719. doi: 10.1016/j.theriogenology.2011.01.008 pmid: 21396699
 15. Rinaudo P, Schultz RM. Effects of embryo culture on global pattern of gene expression in preimplantation mouse embryos. *Reproduction.* 2004;128(3):301-311. doi: 10.1530/rep.1.00297 pmid: 15333781
 16. Gardner DK, Lane M. Ex vivo early embryo development and effects on gene expression and imprinting. *Reprod Fertil Dev.* 2005;17(3):361-370. doi: 10.1071/RD04103 pmid: 15745644
 17. Lane M, Gardner DK. Amino acids and vitamins prevent culture-induced metabolic perturbations and associated loss of viability of mouse blastocysts. *Hum Reprod.* 1998;13(4):991-997. doi: 10.1093/humrep/13.4.991 pmid: 9619560
 18. Koot YE, Macklon NS. Embryo implantation: biology, evaluation, and enhancement. *Curr Opin Obstet Gynecol.* 2013;25(4):274-279. doi: 10.1097/GCO.0b013e3283630d94 pmid: 23787799
 19. Ng EHY, Yeung WSB, Lau EYL, So WWK, Ho PC. High serum oestradiol concentrations in fresh IVF cycles do not impair implantation and pregnancy rates in subsequent frozen-thawed embryo transfer cycles. *Hum Reprod.* 2000;15(2):250-255.
 20. Pollard J, Leibo S. Chilling sensitivity of mammalian embryos. *Theriogenology.* 1994;41(1):101-106.
 21. Duby R, Hill J, O'Callaghan D, Overstrom E, Boland M. Changes induced in the bovine zona pellucida by ovine and bovine oviducts. *Theriogenology.* 1997;47(1).
 22. Slimane W, Heyman Y, Lavergne Y, Humblot P, Renard JP. Assessing chromosomal abnormalities in two-cell bovine *in vitro*-fertilized embryos by using fluorescent *in situ* hybridization with three different cloned probes. *Biol Reprod.* 2000;62(3):628-635. pmid: 10684804
 23. Crosier AE, Farin PW, Dykstra MJ, Alexander JE, Farin CE. Ultrastructural morphometry of bovine blastocysts produced *in vivo* or *in vitro*. *Biol Reprod.* 2001;64(5):1375-1385. doi: 10.1095/biolreprod64.5.1375 pmid: 11319141
 24. Giritharan G, Talbi S, Donjacour A, Di Sebastiano F, Dobson AT, Rinaudo PF. Effect of *in vitro* fertilization on gene expression and development of mouse preimplantation embryos. *Reproduction.* 2007;134(1):63-72. doi: 10.1530/REP-06-0247 pmid: 17641089
 25. Lonergan P, Fair T, Corcoran D, Evans AC. Effect of culture environment on gene expression and developmental characteristics in IVF-derived embryos. *Theriogenology.* 2006;65(1):137-152. doi: 10.1016/j.theriogenology.2005.09.028 pmid: 16289260
 26. Carrasco B, Boada M, Rodriguez I, Coroleu B, Barri PN, Veiga A. Does culture medium influence offspring birth weight? *Fertil Steril.* 2013;100(5):1283-1288. doi: 10.1016/j.fertnstert.2013.07.006 pmid: 23916797
 27. Lane M. Mechanisms for managing cellular and homeostatic stress *in vitro*. *Theriogenology.* 2001;55(1):225-236. doi: 10.1016/S0093-691X(00)00456-8 pmid: 11198084
 28. Kovai B, Vlaisavljevi V. Importance of Blastocyst Morphology in Selection for Transfer. *Advances in embryo transfer: IntechOpen;* 2012. p. 161.

29. Barahona P, Leigh D, Ritchie W, McArthur SJ, Jansen RP. Array CGH and Partial Genome Sequencing for Rapidly Karyotyping IVF Blastocysts Before Single Transfer. Screening the Single Euploid Embryo: Springer; 2015. p. 163-178.
30. Cockburn K, Rossant J. Making the blastocyst: lessons from the mouse. *J Clin Invest.* 2010;**120**(4):995-1003. doi: 10.1172/JCI41229 pmid: 20364097
31. Arnold SJ, Robertson EJ. Making a commitment: cell lineage allocation and axis patterning in the early mouse embryo. *Nat Rev Mol Cell Biol.* 2009;**10**(2):91-103. doi: 10.1038/nrm2618 pmid: 19129791
32. Chazaud C, Yamanaka Y, Pawson T, Rossant J. Early lineage segregation between epiblast and primitive endoderm in mouse blastocysts through the Grb2-MAPK pathway. *Dev Cell.* 2006;**10**(5):615-624. doi: 10.1016/j.devcel.2006.02.020 pmid: 16678776
33. Chen L, Nakai M, Belton RJ, Jr., Nowak RA. Expression of extracellular matrix metalloproteinase inducer and matrix metalloproteinases during mouse embryonic development. *Reproduction.* 2007;**133**(2):405-414. doi: 10.1530/rep.1.01020 pmid: 17307908
34. Dhali A, Anchamparathy VM, Butler SP, Pearson RE, Mullarky IK, Gwazdauskas FC. Gene expression and development of mouse zygotes following droplet vitrification. *Theriogenology.* 2007;**68**(9):1292-1298. doi: 10.1016/j.theriogenology.2007.08.030 pmid: 17915304
35. Elmore S. Apoptosis: a review of programmed cell death. *Toxicol Pathol.* 2007;**35**(4):495-516. doi: 10.1080/01926230701320337 pmid: 17562483
36. Quinn P, Warnes GM, Kerin JF, Kirby C. Culture factors affecting the success rate of *in vitro* fertilization and embryo transfer. *Ann N Y Acad Sci.* 1985;**442**(1):195-204. doi: 10.1111/j.1749-6632.1985.tb37520.x pmid: 3860035
37. Du T, Jia J, Lin X, Xie R, Li J, Xiao D, et al. Generation of Rm21LG transgenic mice: a powerful tool to generate conditional overexpression of miR-21 that is involved in oncogenesis. *Biotechnol Lett.* 2014;**36**(1):9-20. doi: 10.1007/s10529-013-1327-y pmid: 24068500
38. Choi YJ, Gurunathan S, Kim D, Jang HS, Park WJ, Cho SG, et al. Rapamycin ameliorates chitosan nanoparticle-induced developmental defects of preimplantation embryos in mice. *Oncotarget.* 2016;**7**(46):74658-74677. doi: 10.18632/oncotarget.10813 pmid: 27463007
39. Velásquez A, Manríquez J, Castro F, Rodríguez-Alvarez L. Effect of zona pellucida removal on early development of *in vitro* produced bovine embryos. *Arch Med Vet.* 2013;**45**(1).
40. Montag M, Koll B, Holmes P, van der V. Significance of the number of embryonic cells and the state of the zona pellucida for hatching of mouse blastocysts *in vitro* versus *in vivo*. *Biol Reprod.* 2000;**62**(6):1738-1744. doi: 10.1095/biolreprod62.6.1738 pmid: 10819778
41. Hammadeh ME, Fischer-Hammadeh C, Ali KR. Assisted hatching in assisted reproduction: a state of the art. *J Assist Reprod Genet.* 2011;**28**(2):119-128. doi: 10.1007/s10815-010-9495-3 pmid: 21042844
42. Schiewe MC, Hazeleger NL, Scimienti C, Balmaceda JP. Physiological characterization of blastocyst hatching mechanisms by use of a mouse antihatching model*+*Presented in part at the 42nd Annual Meeting of The Pacific Coast Fertility Society, Palm Springs, California, April 21 to 24, 1994.†Supported in part by a Department of Obstetrics and Gynecology Memorial Health Services grant at the University of California Irvine and by resources made available at the University of California Irvine-Beckman Laser Institute. *Fertil Steril.* 1995;**63**(2):288-294. doi: https://doi.org/10.1016/S0015-0282(16)57357-9
43. Staun-Ram E, Goldman S, Gabarin D, Shalev E. Expression and importance of matrix metalloproteinase 2 and 9 (MMP-2 and -9) in human trophoblast invasion. *Reprod Biol Endocrinol.* 2004;**2**(1):59. doi: 10.1186/1477-7827-2-59 pmid: 15294019
44. Peters TJ, Albieri A, Bevilacqua E, Chapman BM, Crane LH, Hamlin GP, et al. Differentiation-dependent expression of gelatinase B/matrix metalloproteinase-9 in trophoblast cells. *Cell Tissue Res.* 1999;**295**(2):287-296. doi: 10.1007/s004410051235 pmid: 9931375
45. Koot YE, Teklenburg G, Salker MS, Brosens JJ, Macklon NS. Molecular aspects of implantation failure. *Biochim Biophys Acta.* 2012;**1822**(12):1943-1950. doi: 10.1016/j.bbadis.2012.05.017 pmid: 22683339
46. Ventura-Junca P, Irarrazaval I, Rolle AJ, Gutierrez JI, Moreno RD, Santos MJ. *In vitro* fertilization (IVF) in mammals: epigenetic and developmental alterations. Scientific and bioethical implications for IVF in humans. *Biol Res.* 2015;**48**(1):68. doi: 10.1186/s40659-015-0059-y pmid: 26683055
47. Kazemi P, Dashtizad M, Shamsara M, Mahdavezhad F, Hashemi E, Fayazi S, et al. Effect of blastocoel fluid reduction before vitrification on gene expression in mouse blastocysts. *Mol Reprod Dev.* 2016;**83**(8):735-742. doi: 10.1002/mrd.22681 pmid: 27409768
48. Henderson GR, Brahmasani SR, Yelisetti UM, Konijeti S, Katari VC, Sisinthy S. Candidate gene expression patterns in rabbit preimplantation embryos developed *in vivo* and *in vitro*. *J Assist Reprod Genet.* 2014;**31**(7):899-911. doi: 10.1007/s10815-014-0233-0 pmid: 24760721
49. Purpera MN, Giraldo AM, Ballard CB, Hylan D, Godke RA, Bondioli KR. Effects of culture medium and protein supplementation on mRNA expression of *in vitro* produced bovine embryos. *Mol Reprod Dev.* 2009;**76**(8):783-793. doi: 10.1002/mrd.21028 pmid: 19288542
50. Ng HH, Surani MA. The transcriptional and signalling networks of pluripotency. *Nat Cell Biol.* 2011;**13**(5):490-496. doi: 10.1038/ncb0511-490 pmid: 21540844
51. Strumpf D, Mao CA, Yamanaka Y, Ralston A, Chawengsaksophak K, Beck F, et al. Cdx2 is required for correct cell fate specification and differentiation of trophectoderm in the mouse blastocyst. *Development.* 2005;**132**(9):2093-2102. doi: 10.1242/dev.01801 pmid: 15788452
52. Lanner F. Lineage specification in the early mouse embryo. *Exp Cell Res.* 2014;**321**(1):32-39. doi: 10.1016/j.yexcr.2013.12.004 pmid: 24333597
53. Niwa H, Toyooka Y, Shimosato D, Strumpf D, Takahashi K, Yagi R, et al. Interaction between Oct3/4 and Cdx2 determines trophectoderm differentiation. *Cell.* 2005;**123**(5):917-929. doi: 10.1016/j.cell.2005.08.040 pmid: 16325584

54. Oron E, Ivanova N. Cell fate regulation in early mammalian development. *Phys Biol*. 2012;**9**(4):045002. doi: [10.1088/1478-3975/9/4/045002](https://doi.org/10.1088/1478-3975/9/4/045002) pmid: 22871593
55. Awonuga AO, Zhong W, Abdallah ME, Slater JA, Zhou SC, Xie YF, et al. Eomesodermin, HAND1, and CSH1 proteins are induced by cellular stress in a stress-activated protein kinase-dependent manner. *Mol Reprod Dev*. 2011;**78**(7):519-528. doi: [10.1002/mrd.21342](https://doi.org/10.1002/mrd.21342) pmid: 21710638
56. Wang Y, Smedberg JL, Cai KQ, Capo-Chichi DC, Xu XX. Ectopic expression of GATA6 bypasses requirement for Grb2 in primitive endoderm formation. *Dev Dyn*. 2011;**240**(3):566-576. doi: [10.1002/dvdy.22447](https://doi.org/10.1002/dvdy.22447) pmid: 20925113
57. Jacobson MD, Weil M, Raff MC. Programmed cell death in animal development. *Cell*. 1997;**88**(3):347-354. pmid: 9039261
58. Hemann MT, Lowe SW. The p53-Bcl-2 connection. *Cell Death Differ*. 2006;**13**(8):1256-1259. doi: [10.1038/sj.cdd.4401962](https://doi.org/10.1038/sj.cdd.4401962) pmid: 16710363